arXiv:1209.4283v2 [math.GT] 16 Oct 2012

The untwisted Dijkgraaf-Witten invariants of links

Haimiao Chen *

School of Mathematical Science, Peking University, Beijing, China

Abstract

In this article we study 3d untwisted Dijkgraaf-Witten theory. We are able to express the DW invariant based on a finite group of a link in terms of its Reshetikhin-Turaev invariants constructed from the quantum double of the same group. Then we develop a feasible method to compute the DW invariants of Montesinos knots. As an application, we derive a formula for the number of n-colorings of Montesinos knots by computing the DW invariant when the finite group is D_n with n odd.

key words: Dijkgraaf-Witten invariant, Reshetikhin-Turaev invariant, extended TQFT, fundamental group, Montesinos knot, number of colorings.

MSC2010: 57M25, 57M27, 57R56

1 Introduction

Dijkgraaf-Witten theory is a topological quantum field theory (TQFT). It was first proposed in [5], and since then has been further studied by many people; see [7–9, 16, 24], and the references therein. It can be defined for any integer $l \geqslant 1$, any finite group Γ and any cohomology class $\alpha \in H^l(\Gamma; U(1))$, and is said to be untwisted when $\alpha = 0$. The key ingredient of the untwisted DW theory (denoted as $\mathrm{DW}^0_{l,\Gamma}$) is counting homomorphisms from the fundamental group of a manifold to Γ.

In dimensional 3, there are a wide class of TQFTs. According to Reshetikhin-Turaev [20], starting from a "modular Hopf algebra", one can firstly associate to each colored framed link (L, c) an invariant $\mathcal{F}(L, c)$, and then use it to construct cobordism invariant, which fits in TQFT axioms. Actually this gives an extended TQFT, see also [2].

From a finite group Γ, a modular Hopf algebra called the (untwisted) quantum double $D(\Gamma)$ (see [2] Section 3.2) can be constructed. So it gives rise to a 3d TQFT, denoted as RT^0_Γ. It was believed [7] that $\mathrm{DW}^0_{3,\Gamma}$ and RT^0_Γ are essentially the same. This was supported by an argument in [1] which concluded that $\mathrm{DW}^0_{3,\Gamma}(M) = \mathrm{RT}^0_\Gamma(M)$ for any closed 3-manifold M.

---

[*] Email: *chenhm@math.pku.edu.cn*

In this article we relate $\mathrm{DW}_{3,\Gamma}^0$ to RT_{Γ}^0 by showing that, for a link $L \subset S^3$, the coefficients of the vector $\mathrm{DW}_{3,\Gamma}^0(S^3 - N(L))$ (where $N(L)$ is a tubular neighborhood of L) with respect to a canonical basis are given by $\frac{1}{\#\Gamma} \cdot \mathcal{F}(L,c)$, for various colorings c. The result is stated precisely as Theorem 4.4.

To reach this goal, it is necessary to consider DW theory as an extended TQFT. This was already done in [8, 9, 16]. The novelty of our work is to build a "visible" bridge connecting $\mathrm{DW}_{3,\Gamma}^0$ to RT_{Γ}^0, so one can see why these two approaches should give rise to the same TQFT. We aim to keep the article on an elementary level, filling in many important details that have not been explained before. Also, when dealing with 3-manifolds with corners, we prefer the concept "$\langle 2 \rangle$-manifold" of [12] to the more widely used "cospan" of [16].

It is hopeful to completely establish that $\mathrm{DW}_{3,\Gamma}^0 = \mathrm{RT}_{\Gamma}^0$ as extended TQFTs. In this article we have, based on $\mathrm{DW}_{3,\Gamma}^0$, associated to each colored framed link (L,c) another invariant $F(L,c)$ and shown $F(L,c) = \mathcal{F}(L,c)$. It is sufficient to show that the cobordism invariant constructed from $F(L,c)$ by Reshetikhin-Turaev's method recovers that of $\mathrm{DW}_{3,\Gamma}^0$. But this requires still much more work. We may do it and generalize to twisted cases in future work.

DW theory not only has theoretical importance since it is one of the first rigorously constructed TQFTs, but also is practically interesting because of its relation with fundamental group. The fundamental group of a manifold, as a noncommutative object, encodes much topological information of the manifold, but at the same time, is often hard to handle. The DW invariant of a manifold, however, can extract partial information of its fundamental group.

In general, enumeration of homomorphisms from a group G to finite groups is usually helpful to understand the group G. For example, the old topic of counting finite-index subgroups of a finitely generated group G is directly related to counting homomorphisms from G to symmetric groups, see [14, 22]; the Hall invariant of a group G, which counts epimorphisms from G onto finite groups, dates back to [10] and also has its own interests [15, 22].

When G is the fundamental group of some manifold M, homomorphisms from G to finite groups have additional topological meanings. In dimension 2, people studied existence and enumeration problems on surface coverings by counting homomorphisms from surface groups to finite groups, see [11, 18, 23]. But in 3-dimension, till now, besides [13] and the author's work [3], there are few such results. In knot theory, there are good reasons that people should pay attention to homomorphisms from $\pi_1(K) := \pi_1(S^3 - N(K))$ to finite groups. It was shown in [6] that all finite quandle invariants can be expressed in terms of, for appropriate finite group Γ and various commuting elements x, y of Γ, the numbers $N_\Gamma(x,y)$ of homomorphisms $\pi_1(K) \to \Gamma$ sending the meridian and the longitude to x and y, respectively; it was also pointed out that such enumeration is helpful to somehow keep track of the knot group since the knot group is residually finite.

For a knot K, $\mathrm{DW}_{3,\Gamma}^0(S^3 - N(K))$ is a vector whose components are exactly $N_\Gamma(x,y)$ in the previous paragraph. We develop a method to compute this invariants for Montesinos knots (see Section 5 for the definition), showing that DW invariants are computable, at least for Montesinos knots. As an illustration, we partially compute the DW invariant when $\Gamma = D_n$ (the dihedral group) with n odd and apply the result to derive a formula for an invariant of long history, namely, the number of n-colorings of knots.

Actually our method is applicable to any other Reshetikhin-Turaev type TQFT. For example, for Montesinos knots, the Witten-Reshetikhin-Turaev invariants which are given by the TQFT constructed from the quantum group $U_q(\mathfrak{g})$ ($\mathfrak{g}$ being a Lie algebra) at roots of unit, can be systematically computed. Finally, the partition function of Reshetikhin-Turaev type TQFT for any closed 3-manifold resulting by doing surgery along a Montesinos knot is computable.

The context is organized as follows. Section 2 is a preliminary on DW theory, tangle and link, and RT theory. Section 3 is devoted to (partially) establishing 3d extended untwisted DW theory. The main result, which expresses the DW invariant of a link in terms of its RT invariants for various colorings, is proved in Section 4. In Section 5 we describe how to compute the DW/RT invariants of Montesinos knots. In Section 6 we derive a formula for the number of n-colorings. The last section is an appendix consisting of the proofs of all important assertions in Section 3; this makes the article more readable.

Notations & conventions.

In this article, all manifolds are compact and oriented. For a manifold M, let $-M$ denote the manifold obtained by reversing the orientation of M.

The torus $S^1 \times S^1$ is denoted as Σ_1. Call $S^1 \times 1$ and $1 \times S^1$ the standard meridian and longitude, respectively.

For elements x, g of a group, denote ${}^{g}x = gxg^{-1}$; for tuples $\mathbf{x} = (x_1, \cdots, x_m)$ and $\mathbf{g} = (g_1, \cdots, g_m)$, abbreviate $({}^{g_1}x_1, \cdots, {}^{g_m}x_m)$ to ${}^{\mathbf{g}}\mathbf{x}$.

All inner product spaces are finite-dimensional; inner product is written to be linear in the first variable and anti-linear in the second.

Let $\mathcal{V}$ be the category of inner product spaces. Equip $\mathbb{C}^m$ with the standard inner product. For $V \in \mathcal{V}$, let $\overline{V} \in \mathcal{V}$ have the same underlying abelian group as V, equipped with the scalar multiplication $(\lambda, v) \mapsto \overline{\lambda}v$, and the inner product $(u, v)_{\overline{V}} = (v, u)_V$. If $\lambda > 0$, then by $\lambda \cdot V$ we mean the inner product space with the same underlying vector space as V and equipped with the inner product $(u, v)_{\lambda \cdot V} = \lambda \cdot (u, v)_V$.

2 Preliminary

2.1 Dijkgraaf-Witten theory

The references for this section are [7–9].

Let l be a positive integer. An *l-dimensional TQFT Z* associates to each closed $(l-1)$-manifold B an inner product space $Z(B)$, and to each l-manifold C an element $Z(C)$ of $Z(\partial C)$, such that the following axioms hold:

(A) (Functoriality) Each diffeomorphism $f : B \to B'$ induces an isometry $f_* : Z(B) \to Z(B')$; for each diffeomorphism $F : C \to C'$, one has $(\partial F)_*(Z(C)) = Z(C')$, where $\partial F : \partial C \to \partial C'$ is the induced map on boundaries.

(B) (Orientation) There are natural isometries $Z(-B) \cong \overline{Z(B)}$, through which $Z(-C)$ is sent to $Z(C)$.

(C) (Multiplicativity) There are natural isometries $Z(B \sqcup B') \cong Z(B) \otimes Z(B')$, through which $Z(C \sqcup C')$ is sent to $Z(C) \otimes Z(C')$.

(D) (Gluing) If $\partial C_1 = B_1 \sqcup B_3$ and $\partial C_2 = -B_3 \sqcup B_2$, one can glue C_1 and C_2 along B_3 to obtain a new manifold C_3. Then $Z(C_3) = \langle Z(C_1), Z(C_2)\rangle$, where $\langle \cdot, \cdot \rangle$ is the pairing

$$Z(B_1) \otimes Z(B_3) \otimes \overline{Z(B_3)} \otimes Z(B_2) \to Z(B_1) \otimes Z(B_2).$$

(E) (Normalization) $Z(\emptyset^{l-1}) = \mathbb{C}$, $Z(\emptyset^l) = 1$, where $\emptyset^{l-1}$ and $\emptyset^l$ are the empty set viewed as $(l-1)$- and l-dimensional manifolds, respectively.

The l-dimensional untwisted DW theory Z based on a finite group Γ is defined as follows.

For a connected closed $(l-1)$-manifold B, let

$$Z(B) = \frac{1}{\#\Gamma} \cdot \Big(\bigoplus_{\phi : \pi_1(B) \to \Gamma} \mathbb{C}\phi \Big)^{\Gamma}; \tag{1}$$

for a connected closed l-manifold C, let

$$Z(C) = \frac{1}{\#\Gamma} \cdot \#\{\Phi : \pi_1(C) \to \Gamma\}; \tag{2}$$

for a connected l-manifold C with nonempty boundary, let

$$Z(C)(\phi) = \#\{\Phi : \pi_1(C) \to \Gamma \colon \Phi \circ \iota_* = \phi\}, \tag{3}$$

where $\iota_* : \pi_1(\partial C) \to \pi_1(C)$ is induced by the inclusion $\iota : \partial C \hookrightarrow C$.

When $l = 3$. It is easy to get

$$Z(S^2) = \frac{1}{\#\Gamma} \cdot \mathbb{C}, \qquad Z(D^3) = 1 \in Z(S^2). \tag{4}$$

The inner product space $E := Z(\Sigma_1)$ can be identified with the vector space of functions $f : \{(x, g) \in \Gamma^2 \colon xg = gx\} \to \mathbb{C}$ such that $f({}^a x, {}^a g) = f(x, g)$ for all $a \in \Gamma$, with the inner product given by

$$(f_1, f_2) = \frac{1}{\#\Gamma} \cdot \sum_{xg = gx} f_1(x, g)\overline{f_2(x, g)}. \tag{5}$$

For each conjugacy class c of Γ, take $x \in c$, choose an irreducible character ρ of the centralizer $C(x)$ of x, and let $\chi_{c,\rho}({}^g x, {}^g h) = \rho(h)$ for $h \in C(x)$ and $\chi_{c,\rho}(y, k) = 0$ whenever $y \notin c$. It is known (see Lemma 5.4 of [9]) that E has a canonical orthonormal basis consisting of all such $\chi_{c,\rho}$.

The mapping class group of Σ_1 is isomorphic to $\mathrm{SL}(2,\mathbb{Z})$ and acts on E by

$$\left(\begin{pmatrix} a & b \\ c & d \end{pmatrix}_* f\right)(x,g) = f(x^a g^b, x^c g^d). \tag{6}$$

In particular, for $Q = \begin{pmatrix} 1 & 0 \\ 1 & 1 \end{pmatrix}$, the expression is

$$Q_* \chi_{c,\rho} = \theta_{c,\rho} \cdot \chi_{c,\rho}, \tag{7}$$

where

$$\theta_{c,\rho} = \chi_{c,\rho}(x,x)/\chi_{c,\rho}(x,e) \text{ for an arbitrary } x \in c. \tag{8}$$

2.2 Framed Tangles and links

A *framed tangle* T is an equivalence class of pairs (A, fr) where A is a 1-manifold embedded in $[0,1] \times \mathbb{R}^2 \subset \mathbb{R}^3$ such that $\partial A \subset \{0,1\} \times \mathbb{R} \times \{0\}$, and fr is a normal vector field on A. Two pairs (A, fr) and $(A', \mathrm{fr'})$ are equivalent if there is an orientation-preserving diffeomorphism of $\mathbb{R}^3$ that takes A to A' and takes fr to fr'. Denote $T = [A, \mathrm{fr}]$. Below we sometimes do not distinguish a framed tangle from its representatives. Call $(\{0\} \times \mathbb{R} \times \{0\}) \cap A$ (resp. $(\{1\} \times \mathbb{R} \times \{0\}) \cap A$) the *source* (resp. the *target*) of T and denote it as $s(T)$ (resp. $t(T)$). Note that $s(T)$ and $t(T)$ should be considered as oriented 0-manifolds.

Tangles with empty source and target are just links.



Figure 1: The closure of a tangle

A framed tangle can always be presented as a tangle diagram, with the blackboard framing understood. Hence two tangle diagrams represent the same framed tangle if and only if they represent the same unframed tangle and the writhe numbers of corresponding components are the same. When drawing a tangle, we always put its target on the top and the source on the bottom, so that tangles are read from down to up.



Figure 2: Three of the basic tangles



Figure 3: Integral tangles

Each framed tangle T can be uniquely decomposed into the union of four framed sub-tangles: $T = T^0 \sqcup T^- \sqcup T^+ \sqcup T^1$, such that $s(T^0) = t(T^0) = t(T^-) = s(T^+) = \emptyset$, and $s(T^-), t(T^+), s(T^1), t(T^1) \neq \emptyset$.

Given two tangles T_1, T_2, the *horizontal composite* $T_1 \sqcup T_2$ is defined by putting T_2 on the right side of T_1. When $\#s(T_2) = \#t(T_1)$ (so there is an obvious bijection between $s(T_2)$ and $t(T_1)$), and the orientations are compatible, write $s(T_2) = t(T_1)$, and define the *vertical composite* $T_2 \cdot T_1$ to be framed tangle obtained by identifying $t(T_1)$ with $s(T_2)$ in the obvious way, so that $s(T_2 \cdot T_1) = s(T_1)$ and $t(T_2 \cdot T_1) = t(T_2)$.

When $s(T) = t(T)$, the *closure* of T, denoted $\overline{T}$, is the closed 1-manifold obtained by connecting the source and target using parallel lines in the way shown in Figure 1.

Call the tangles in Figure 2 and the $\pm\mathbf{1}$ in Figure 3 *basic tangles*.

For $n \in \mathbb{Z}$, the *integral tangle* $\mathbf{n}$ is the iterated vertical composite of $|n|$ copies of $\mathbf{1}/\mathbf{-1}$ when n is positive/negative.

Definition 2.1. *For a link $L \subset S^3$, the* DW invariant of L, *denoted as $Z(L)$, is the image of $Z(S^3 - N(L))$ under the isometry $Z(\partial N(L)) \cong E^{\otimes \#\pi_0(L)}$ induced by the diffeomorphism $\partial N(L) \to \sqcup_{\#\pi_0(L)} \Sigma_1$.*

Remark 2.2. *For each component K of a framed link L, choose a longitude $l(K)$ according to the framing and a meridian $m(K)$, then the isomorphism $\pi_1(\partial N(K)) \cong \pi_1(\Sigma_1)$ induced by the diffeomorphism $\partial N(K) \to \Sigma_1$ sends $[m(K)]$ to $[m_0]$ and sends $[l(K)]$ to $[m_0]^{wr(K)}[l_0]$, where m_0/l_0 is the standard meridian/longitude of Σ_1, and $wr(K)$ is the writhe number of K.*

Suppose $L = K_1 \sqcup \cdots \sqcup K_r$, *then*

$$\begin{aligned}&\#\{\Phi : \pi_1(S^3 - N(L)) \to \Gamma \colon \Phi([m(K_i)]) = x_i, \Phi([l(K_i)]) = g_i, 1 \leqslant i \leqslant r\}\\ =&((Q_*^{wr(K_1)} \otimes \cdots \otimes Q_*^{wr(K_r)})Z(L))(x_1, g_1; \cdots ; x_r, g_r).\end{aligned}$$

2.3 The category $\mathcal{E}$

Fix a finite group Γ. Let $\mathcal{E}$ be the category of graded inner product spaces $U = \bigoplus_{x\in\Gamma} U_x$ together with a left unitary Γ action such that $g(U_x) = U_{{}^g x}$; a morphism $U \to V$ is a family of linear maps $f = \{f_x : U_x \to V_x \colon x \in \Gamma\}$ such that $g \circ f_x = f_{{}^g x} \circ g$ for all x, g.

Suppose $U \in \mathcal{E}$, and $xg = gx$, let

$$\chi_U(x, g) = \mathrm{tr}(g : U_x \to U_x). \tag{9}$$

The function χ_U belongs to E and is called the *character* of U. Define

$$\mathrm{Dim} U = \sum_{x\in\Gamma} \chi_U(x, e). \tag{10}$$

As is well-known [2], the category $\mathcal{E}$ is just the category of finite-dimensional modules of the quantum double $D(\Gamma)$, and it is a so-called “modular tensor category”. It has finitely many isomorphism classes of simple objects; let Λ denote the set of them. Actually $\{\chi_U \colon U \in \Lambda\}$ is the same as the canonical orthonormal basis $\{\chi_{c,\rho}\}$ mentioned in Section 3.1. There are some crucial structures of $\mathcal{E}$:

- A bifunctor called *tensor product*

  $$\begin{aligned}\odot : \mathcal{E} \times \mathcal{E} \to& \mathcal{E},\\ (U \odot V)_x =& \bigoplus_{x_1 x_2 = x} U_{x_1} \otimes V_{x_2},\end{aligned} \tag{11}$$

  with a “unit” object $\underline{\mathbb{C}}$ with

  $$\underline{\mathbb{C}}_e = \mathbb{C}, \quad \underline{\mathbb{C}}_x = 0 \text{ for all } x \neq e. \tag{12}$$

- A natural isometry called *associator*

  $$(U \odot V) \odot W \to U \odot (V \odot W) \tag{13}$$

  given by the natural isomorphism of vector spaces

  $$(U_x \otimes V_y) \otimes W_z \cong U_x \otimes (V_y \otimes W_z). \tag{14}$$

- A natural isometry called *braiding*

  $$R_{U,V} : U \odot V \to V \odot U, \tag{15}$$

  $$u_x \otimes v_y \mapsto (x v_y)_{{}^x y} \otimes u_x, \quad \text{for } u_x \in U_x, v_y \in V_y. \tag{16}$$

- An involution called *dual*

$$^{\star}:\mathcal{E}\to\mathcal{E},\qquad (U^{\star})_x=\overline{U_{x^{-1}}}; \tag{17}$$

the action $g:(U^{\star})_x\to(U^{\star})_{{}^g x}$ is the same as a map as $g:U_{x^{-1}}\to U_{{}^g x^{-1}}$.

- A pair of natural transformations

$$i_U:\underline{\mathbb{C}}\to U\odot U^{\star},\qquad e_U:U^{\star}\odot U\to\underline{\mathbb{C}}, \tag{18}$$

which are expressed explicitly as

$$i_U(1)=\sum_x\sum_a u_x^a\otimes\overline{u_x^a}, \tag{19}$$

$$e_U(\overline{u_x}\otimes u'_y)=\delta_{x,y}\cdot(u'_y,u_x)_{U_x},\qquad \text{for } u_x\in U_x,u'_y\in U_y, \tag{20}$$

where $\{u_x^a\}$ is an arbitrary orthonormal basis of U_x, and $\overline{u_x}$ is the image of u_x under the identification $\overline{U_x}=U^{\star}_{x^{-1}}$.

For $m\geqslant 1$, let $\mathcal{E}^{\boxtimes m}$ be the category of graded inner product spaces $U=\bigoplus_{\mathbf{x}\in\Gamma^m}U_{\mathbf{x}}$ together with a left unitary Γ^m action such that $\mathbf{g}U_{\mathbf{x}}=U_{{}^{\mathbf{g}}\mathbf{x}}$. There is a "2-inner product" on $\mathcal{E}$ which is given by, for $U=\bigoplus_{\mathbf{x}}U_{\mathbf{x}}$, $V=\bigoplus_{\mathbf{x}}V_{\mathbf{x}}$,

$$(U,V)_{\mathcal{E}^{\boxtimes m}}=\frac{1}{(\#\Gamma)^m}\cdot\Big(\bigoplus_{\mathbf{x}\in\Gamma^m}U_{\mathbf{x}}\otimes\overline{V_{\mathbf{x}}}\Big)^{\Gamma^m}. \tag{21}$$

We have a natural isomorphism

$$(U,V)_{\mathcal{E}^{\boxtimes m}}\cong\mathcal{E}^{\boxtimes m}(V,U). \tag{22}$$

For $U(1),\cdots,U(m)\in\mathcal{E}$, let $U=U(1)\boxtimes\cdots\boxtimes U(m)$ with $U_{\mathbf{x}}=\bigotimes_{i=1}^{m}U(i)_{x_i}$, and call U the *external tensor product* of $U(1),\cdots,U(m)$.

The functor $\mathcal{E}\boxtimes\mathcal{E}\to\mathcal{E}$ given by sending $U\boxtimes V$ to $U\odot V$ and extending "linearly" to the whole domain is well-defined; we denote it also by $\odot$.

2.4 Reshetikhin-Turaev invariants

Definition 2.3. *A colored framed tangle is a pair (T,c) consisting of a framed tangle T and a map $c:\pi_0(T)\to\Lambda$.*

Two colored framed tangles (T,c) and (T',c') are considered as the same if T' is obtained by reversing the orientations of some components $A_i,i\in I$, of T and c' differs from c only at these components with $c'(A_i)=c(A_i)^{\star},i\in I$.

Given a colored framed tangle (T,c). If $s(T)=\emptyset$ (resp. $t(T)=\emptyset$), then put $U_T=\underline{\mathbb{C}}$ (resp. $U^T=\underline{\mathbb{C}}$). Otherwise, suppose $s(T)=(s_1,\cdots,s_k),t(T)=$



Figure 4: Colored basic tangles

$(t_1, \cdots, t_l)$. Define $U_{(i)}$ to be $c(A)$ (resp. $c(A)^\star$) if s_i is the initial (resp. terminal) point of some component A of T, and define $U^{(j)}$ to be $c(A)^\star$ (resp. $c(A)$) if t_j is the initial (resp. terminal) point of some component A of T; Put

$$U_T = U_{(1)} \odot \cdots \odot U_{(k)}, \qquad U^T = U^{(1)} \odot \cdots \odot U^{(l)}. \tag{23}$$

As done in [2,20], associated to (T, c) is a morphism

$$\mathcal{F}(T, c) : U_T \to U^T. \tag{24}$$

First, to each colored basic tangles as illustrated in Figure 4, associate a morphism as follows, respectively: (a) id_U; (b) $R_{U,V}$; (c) $R^{-1}_{V,U}$; (d) i_U; (e) e_U.

Second, call a framed tangle *elementary* if it is a horizontal composite of basic tangles; for a colored elementary framed tangle (T, c), let $\mathcal{F}(T, c)$ be the tensor product of the morphisms associated to the basic pieces.

Third, for a general colored framed tangle T, it is always possible to decompose T as a vertical composite of elementary pieces, then set $\mathcal{F}(T, c)$ to be the composite of the morphisms associated to the elementary pieces.

Theorem 2.4. *(Reshetikhin-Turaev) The morphism $\mathcal{F}(T, c)$ is an invariant of colored framed tangles.*

3 3D extended DW theory

The TQFT axioms, especially the gluing axiom, which 3d DW theory satisfies, enable us to compute the DW invariant of a 3-manifold M by cutting M along surfaces into relatively simple pieces and combine the DW invariants of these pieces into a global invariant. Nevertheless, we cannot go too far. For example, if not impossible, at least no one knows how to cut the complement of a knot in S^3 into simple sub-manifolds. So the effect is very limited.

But the situation will be quite different if we allow the pieces to be manifolds "with corners": there will be very few building blocks for 3-manifolds, as we shall see. This motives us to extend DW theory.

Let S^1 denote $\{z \in \mathbb{C}\colon |z| = 1\}$ with the standard orientation and base-point 1. For any 1-manifold, we assume that each connected component A is equipped with a diffeomorphism $A \to S^1$ called *label*, so that A has an induced base-point; we can also use A to denote the loop determined by the orientation and the base-point. For each surface with nonempty boundary, each connected component of its boundary has a label and a basepoint.

For a topological space X, let $\Pi_1(X)$ denote the fundamental groupoid of X. A finite group Γ is regarded as a groupoid with a single object.

Definition 3.1. *The* Γ-fundamental category *of a topological space X, denoted as $\mathfrak{B}^\Gamma(X)$, is the category whose objects are functors $\rho : \Pi_1(X) \to \Gamma$, and morphisms $\eta : \rho \to \rho'$ are maps $\eta : X \to \Gamma$ such that $\rho'([\gamma]) = \eta(t[\gamma])\rho([\gamma])\eta(s[\gamma])^{-1}$ for all $[\gamma]$, where $s[\gamma], t[\gamma]$ are the initial and terminal points of any $\gamma \in [\gamma]$, respectively.*

Call $\rho([\gamma])$ the holonomy *of ρ along $[\gamma]$. Whenever $\eta : \rho \to \rho'$ is a morphism, write $\eta \cdot \rho = \rho'$.*

Remark 3.2. *Here $\mathfrak{B}^\Gamma(X)$ is the same thing as $\mathcal{A}_0(X)_\Gamma$ defined in [16].*

If X is compact, as we always assume to be, then $\mathfrak{B}^\Gamma(X)$ has finitely many connected components.

Suppose $\mathbb{S}$ is a CW structure on X together with a total ordering of 0-cells. Let $\Pi_1(X; \mathbb{S})$ be the full subcategory of $\Pi_1(X)$ having objects the 0-cells of $\mathbb{S}$. Let $F_\mathbb{S} : \Pi_1(X; \mathbb{S}) \to \Pi_1(X)$ be the inclusion functor, and define a functor $G_\mathbb{S} : \Pi_1(X) \to \Pi_1(X; \mathbb{S})$ as follows. For each $x \in X$, let $\sigma(x)$ be the unique cell of lowest dimension that contains x, so $x = \sigma(x)$ if $x \in \mathbb{S}^0$ and $x \in \mathrm{int}(\sigma(x))$ otherwise. Among the vertices of $\sigma(x)$ let $G_\mathbb{S}(x)$ be the first one according to the ordering, and let $P_\mathbb{S}(x)$ be a path in $x \cup G_\mathbb{S}(x) \cup \mathrm{int}(\sigma(x))$ from x to $G_\mathbb{S}(x)$, which is unique up to homotopy since $\mathrm{int}(\sigma(x))$ is contractible. For any morphism $[\gamma] : x \to y$, let $G_\mathbb{S}[\gamma] = [P_\mathbb{S}(y) \cdot \gamma \cdot P_\mathbb{S}(x)^{-1}]$. It is easy to see that $G_\mathbb{S} \circ F_\mathbb{S} = \mathrm{id}$ and $F_\mathbb{S} \circ G_\mathbb{S}$ is naturally isomorphic to id.

Let $\mathfrak{B}^\Gamma(X; \mathbb{S})$ be the category whose objects are morphisms $\varrho : \Pi_1(X; \mathbb{S}) \to \Gamma$ and morphisms are defined similarly as those of $\mathfrak{B}^\Gamma(X)$, then it is equivalent to $\mathfrak{B}^\Gamma(X)$ via pre-composing with the functors $F_\mathbb{S}$ and $G_\mathbb{S}$.

3.1 DW invariants of 2-manifolds

For a connected closed surface B we re-define

$$Z(B) = \bigoplus_{[\rho] \in \pi_0(\mathfrak{B}^\Gamma(X))} \frac{1}{\#\mathrm{Aut}\rho} \cdot \mathbb{C}[\rho], \tag{25}$$

where $\mathbb{C}[\rho] = \mathbb{C}$ and ρ is an arbitrary representative of $[\rho]$. It is easy to see that this definition coincides with (1).

Now suppose B is connected and $\partial B = \sqcup_{i=1}^{p} \partial_i^- B \sqcup_{j=1}^{q} \partial_j^+ B \neq \emptyset$, with $\partial_i^- B = -S^1, \partial_j^+ B = S^1$; by this we mean the labels $\partial_i^- B \to S^1$ are orientation-reversing, while the ones $\partial_j^+ B \to S^1$ are orientation-preserving. We shall define $Z(B)$ as an object of $\mathcal{E}^{\boxtimes(p+q)}$.

For $\mathbf{x} \in \Gamma^p$ and $\mathbf{y} \in \Gamma^q$, let $\mathfrak{B}^{\Gamma}_{\mathbf{y},\mathbf{x}}(B)$ be the subcategory of $\mathfrak{B}^{\Gamma}(B)$ whose objects are functors $\phi : \Pi_1(B) \to \Gamma$ with $\phi(\partial_i^- B) = x_i, \varphi(\partial_j^+ B) = y_j$, and morphisms are maps $\eta : \phi \to \phi'$ with $\eta|_{\partial B} \equiv e$. Define

$$Z(B)_{\mathbf{y},\mathbf{x}} = \bigoplus_{[\rho]\in\pi_0(\mathfrak{B}^{\Gamma}_{\mathbf{y},\mathbf{x}}(B))} \mathbb{C}[\rho]. \tag{26}$$

Put

$$Z(B) = \bigoplus_{\mathbf{x}\in\Gamma^p,\mathbf{y}\in\Gamma^q} Z(B)_{\mathbf{y},\mathbf{x}}; \tag{27}$$

define the action of $\Gamma^p \times \Gamma^q$ on $Z(B)$ by letting $(\mathbf{g},\mathbf{h})$ send $[\rho]$ to

$$\eta_{\mathbf{g},\mathbf{h}} \cdot [\rho] := [\eta_{\mathbf{g},\mathbf{h}} \cdot \rho], \tag{28}$$

where $\eta_{\mathbf{g},\mathbf{h}} : B \to \Gamma$ takes g_i on $\partial_i^- B$, h_j on $\partial_j^+ B$ and e elsewhere.

Remark 3.3. *If $p = 0$, then we write $Z(B)_{\mathbf{y},-}$ instead of $Z(B)_{\mathbf{y},\mathbf{x}}$; similarly for the case when $q = 0$.*

The DW invariant of B can be alternatively defined as a functor $\hat{Z}(B) : \mathcal{E}^{\boxtimes p} \to \mathcal{E}^{\boxtimes q}$; for $U = \bigoplus_{\mathbf{x}\in\Gamma^p} U_{\mathbf{x}}$ and $\mathbf{y} \in \Gamma^q$,

$$(\hat{Z}(B)(U))_{\mathbf{y}} = \frac{1}{(\#\Gamma)^p} \cdot (\bigoplus_{\mathbf{x}} Z(B)_{\mathbf{y},\mathbf{x}^{-1}} \otimes U_{\mathbf{x}})^{\Gamma^p}. \tag{29}$$

For a general surface B, let $Z(B)$ ($\hat{Z}(B)$ resp.) be the external tensor product of $Z(B_i)$ ($\hat{Z}(B_i)$ resp.), taking over all connected components B_i.

Remark 3.4. *Suppose $\mathbb{S}$ is a CW-structure on B such that $\mathbb{S}^0$ contains all the base-points of B. We can define $Z(B;\mathbb{S})$ and $\hat{Z}(B;\mathbb{S})$ similarly as above, with $\mathfrak{B}^{\Gamma}(B)$ replaced by $\mathfrak{B}^{\Gamma}(B;\mathbb{S})$. Then $Z(B) \cong Z(B;\mathbb{S})$ and $\hat{Z}(B)$ is naturally isomorphic to $\hat{Z}(B;\mathbb{S})$. The point is that $Z(B)$ and $\hat{Z}(B)$ are intrinsically defined but hard to view clearly, while $Z(B;\mathbb{S})$ and $\hat{Z}(B;\mathbb{S})$ depend on $\mathbb{S}$ but are convenient to describe. From now on we do not distinguish $Z(B)$ with $Z(B;\mathbb{S})$.*

The DW invariants of 2-manifolds satisfy the following axioms.

Theorem 3.5. *(a) (Functorality) Each diffeomorphism preserving orientations and labels $f : B \to B'$ induces an isometry $f_* : Z(B) \cong Z(B')$ in $\mathcal{E}^{\boxtimes\pi_0(\partial B)}$ and a natural isomorphism $f_* : \hat{Z}(B) \cong \hat{Z}(B')$.*

(b) (Multiplicativity) There is are a natural isometry $Z(B \sqcup B') \cong Z(B) \boxtimes Z(B')$ and a natural isomorphism $\hat{Z}(B \sqcup B') \cong \hat{Z}(B) \boxtimes \hat{Z}(B')$.

(c) (Gluing) Suppose $A \hookrightarrow B$ is a closed 1-dimensional submanifold. Cut B along A into B^{cut}, so that $\partial B^{cut} = -A \sqcup A \sqcup \partial B$. Then there is a natural isometry

$$Z(B) \cong gl_A(Z(B^{cut})),$$

where gl_A is the composite

$$\mathcal{E}^{\boxtimes \pi_0(\partial B^{cut})} \cong \mathcal{E}^{\boxtimes \pi_0(\partial B)} \boxtimes \mathcal{E}^{\boxtimes \pi_0(A)} \boxtimes \mathcal{E}^{\boxtimes \pi_0(A)} \to \mathcal{E}^{\boxtimes \pi_0(\partial B)},$$

$$gl_A(U)_{\mathbf{y}} = \frac{1}{(\#\Gamma)^{\#\pi_0(A)}} \cdot \Big(\bigoplus_{\mathbf{x} \in \Gamma^{\pi_0(A)}} U_{\mathbf{y},\mathbf{x};\mathbf{x}^{-1}} \Big)^{\Gamma^{\pi_0(A)}}. \tag{30}$$

In particular, if B_1, B_2 are surfaces with $\partial^+ B_1 = -\partial^- B_2$, and $B = B_1 \cup_{\partial^+ B_1} B_2$, then there is are a natural isometry $Z(B) \cong gl_{\partial^+ B_1}(Z(B_2) \boxtimes Z(B_1))$ and a natural isomorphism $\hat{Z}(B) \cong \hat{Z}(B_2) \circ \hat{Z}(B_1)$.

Remark 3.6. *In (a) and below, for a set J, we use $\mathcal{E}^{\boxtimes J}$ to denote the functor category $\mathcal{F}un(J, \mathcal{E})$ where J is regarded as a discrete category. Whenever a bijection $J \cong \{1, \cdots, \#J\}$ is chosen, $\mathcal{E}^{\boxtimes J}$ can be identified with $\mathcal{E}^{\boxtimes \#J}$.*

Remark 3.7. *Here we do not discuss orientation axiom, since it is not used in this article. For the same reason, orientation axiom for 3-manifolds is not included.*

3.2 Examples I

(1) Let $D = D^2$ be the 2-dimensional disk. Clearly

$$\hat{Z}(D) : \mathcal{V} \to \mathcal{E}, \qquad W \mapsto \overline{W}, \tag{31}$$

with

$$\overline{W}_e = W, \qquad \overline{W}_x = 0 \text{ for all } x \neq e. \tag{32}$$

(2) Let AN be the annulus, with CW structures shown in Figure 5(a). By definition,

$$Z(AN) = \bigoplus_{x,y} Z(AN)_{y,x} = \bigoplus_{x,y} \Big(\bigoplus_{g :\, y = {}^g x^{-1}} \mathbb{C}\langle x, g \rangle \Big).$$

$$\hat{Z}(AN) : \mathcal{E} \to \mathcal{E}, \qquad (\hat{Z}(AN)(U))_y = \frac{1}{\#\Gamma} \cdot \Big(\bigoplus_{x,g :\, y = {}^g x} \mathbb{C}\langle x^{-1}, g \rangle \otimes U_x \Big)^{\Gamma}.$$

There is an isometry

$$K_U : \hat{Z}(AN)(U) \cong U, \qquad \sum_{x,g :\, y = {}^g x} \langle x^{-1}, g \rangle \otimes u_x^g \mapsto u_y^e, \tag{33}$$

for all y, with inverse

$$K_U^{-1} : U \cong \hat{Z}(AN)(U), \qquad u_y \mapsto \sum_g \langle {}^g y^{-1}, g^{-1} \rangle \otimes g u_y, \tag{34}$$

for all $u_y \in U_y$ and all y. The isometry K_U is natural in U, hence gives rise to a natural isomorphism $K : \hat{Z}(AN) \cong \mathrm{id}$.



Figure 5: (a) The annulus AN; (b) The pair of pants P

(3) Let P be the pair of pants shown in Figure 5(b).

$$Z(P) = \bigoplus_{x, x_1, x_2} Z(P)_{x,(x_1,x_2)} = \bigoplus_{x,(x_1,x_2)} \left(\bigoplus_{\substack{g_1, g_2 : \\ {}^{g_1}x_1^{-1} \cdot {}^{g_2}x_2^{-1} = x}} \mathbb{C}\langle x_1, x_2, g_1, g_2 \rangle \right),$$

$$\begin{aligned} \hat{Z}(P)(U \boxtimes V)_x &= \frac{1}{(\#\Gamma)^2} \cdot \Big(\bigoplus_{\substack{x_1, x_2, g_1, g_2 : \\ {}^{g_1}x_1 \cdot {}^{g_2}x_2 = x}} \mathbb{C}\langle x_1^{-1}, x_2^{-1}, g_1, g_2 \rangle \otimes U_{x_1} \otimes V_{x_2} \Big)^{\Gamma^2} \\ &\cong \bigoplus_{x_1, x_2 :\, x_1 x_2 = x} U_{x_1} \otimes V_{x_2}, \end{aligned}$$

the isometry given by

$$\sum_{\substack{x_1, x_2, g_1, g_2 : \\ {}^{g_1}x_1 \cdot {}^{g_2}x_2 = x}} \langle x_1^{-1}, x_2^{-1}, g_1, g_2 \rangle \otimes u_{x_1,x_2}^{g_1,g_2} \otimes v_{x_1,x_2}^{g_1,g_2} \mapsto \sum_{x_1, x_2 :\, x_1 x_2 = x} u_{x_1,x_2}^{e,e} \otimes v_{x_1,x_2}^{e,e}. \tag{35}$$

Thus $\hat{Z}(P)$ is naturally isomorphic to $\odot : \mathcal{E} \boxtimes \mathcal{E} \to \mathcal{E}$.

Let $CL, B(\cap), B(\cup)$ be the surfaces shown in Figure 6. They are diffeomorphic to each other as manifolds with boundaries, but are pairwise different as cobordisms. We orient and give CW structures to them as the figure illustrates.



Figure 6: (a) CL; (b) $B(\cap)$; (c) $B(\cup)$

(4) Note that CL is the same as AN, so $\hat{Z}(CN) \cong \mathrm{id}$.

(5) Since

$$Z(B(\cap))_{-,(y,x)} = \bigoplus_{y={}^g x^{-1}} \mathbb{C}\langle x, g\rangle,$$

we have, for $U = \bigoplus\limits_{x,y\in\Gamma} U_{y,x} \in \mathcal{E} \boxtimes \mathcal{E}$,

$$\begin{aligned}\hat{Z}(B(\cap))(U) &= \frac{1}{(\#\Gamma)^2} \cdot (\bigoplus_{x,y,g\colon y={}^g x^{-1}} \mathbb{C}\langle x^{-1}, g\rangle \otimes U_{y,x})^{\Gamma^2} \\ &\cong \frac{1}{\#\Gamma} \cdot (\bigoplus_{x\in\Gamma} U_{x^{-1},x})^{\Gamma}; \end{aligned} \tag{36}$$

alternatively,

$$\hat{Z}(B(\cap))(U \boxtimes V) \cong \frac{1}{\#\Gamma} \cdot (\bigoplus_{x} U_{x^{-1}} \otimes V_x)^{\Gamma} = (V, U^{\star})_{\mathcal{E}} \cong \mathcal{E}(U^{\star}, V) \tag{37}$$

the isomorphism given by

$$\sum_{x,i} u_{x^{-1},i} \otimes v_{x,i} \mapsto (\overline{u} \mapsto \sum_{x,i} (u_{x^{-1},i}, u)_U \cdot v_{x,i}). \tag{38}$$

Hence $\hat{Z}(B(\cap)) : \mathcal{E} \boxtimes \mathcal{E} \to \mathcal{V}$ is naturally isomorphic to the functor $U \boxtimes V \mapsto \mathcal{E}(U^{\star}, V)$.

(6) Similarly,

$$Z(B(\cup)) = \bigoplus_{x,y} Z(B(\cup))_{(y,x),-} = \bigoplus_{x,y} \left(\bigoplus_{g\colon y={}^g x^{-1}} \mathbb{C}\langle x, g\rangle \right), \tag{39}$$

and $\hat{Z}(B(\cup)) : \mathcal{V} \to \mathcal{E} \boxtimes \mathcal{E}$ is isomorphic to the functor $W \mapsto W \otimes Z(B(\cup))$.

Denote

$$H(y,x) = \bigoplus_{g\colon y={}^g x^{-1}} \mathbb{C}\langle x, g\rangle, \qquad H = \bigoplus_{x,y} H(y,x) = Z(B(\cup)). \tag{40}$$

There are a natural isomorphism

$$\mathcal{E}^{\boxtimes 2}(H, U \boxtimes V) \cong \mathcal{E}(V^\star, U), \qquad f \mapsto (\overline{v_{x^{-1}}} \mapsto (f\langle x^{-1}, e\rangle, v_{x^{-1}})_V), \tag{41}$$

where $(f\langle x^{-1}, e\rangle, v_{x^{-1}})_V) = \sum_a (v^a, v_{x^{-1}}) \cdot u^a$ if $f\langle x^{-1}, e\rangle = \sum_a u^a \otimes v^a$, and an isometry

$$\hat{Z}(B(\cap))(H) \cong \frac{1}{\#\Gamma}(\bigoplus_{y=x^{-1}} H(y,x))^\Gamma = \frac{1}{\#\Gamma}(\bigoplus_{xg=gx} \mathbb{C}\langle x, g\rangle)^\Gamma = E, . \tag{42}$$

(7) Given a framed tangle T, let $B(T) = T \times S^1$, then $\hat{Z}(B(T)) : \mathcal{E}^{\boxtimes s(T)} \to \mathcal{E}^{\boxtimes t(T)}$ is naturally isomorphic to $\hat{Z}^0(T) \otimes \hat{Z}'(T)$, with

$$\hat{Z}^0(T) : \mathcal{V} \to \mathcal{V}, \qquad W \mapsto W \otimes E^{\otimes \pi_0(T^0)}; \tag{43}$$

and

$$\begin{aligned} &\hat{Z}'(T) : \mathcal{E}^{\boxtimes s(T)} \to \mathcal{E}^{\boxtimes t(T)}, \qquad \boxtimes_{i \in s(T)} U(i) \mapsto V = \bigoplus_{\mathbf{y} \in \Gamma^{t(T)}} V_{\mathbf{y}}, \\ &V_{\mathbf{y}} = \bigotimes_{i \rightharpoondown j} \mathcal{E}(U(j)^\star, U(i)) \bigotimes_{l \in t(T^1)} U(\sigma(l))_{y_l} \bigotimes_{i \rightharpoonup j} H(y_j, y_i), \end{aligned} \tag{44}$$

where $\sigma : t(T^1) \to s(T^1)$ is the bijection induced by T^1, and $i \rightharpoondown j$ ($i \rightharpoonup j$ resp.) means there is a component A of T^- (T^+ resp.) with initial point i and terminal point j. This is obtained by Example (4)-(6) and Theorem 3.5(b).

Now we have an isomorphism

$$\begin{aligned} &\mathcal{E}^{\boxtimes t(T)}(\hat{Z}(B(T))(\boxtimes_{i \in s(T)} U_{(i)}), \boxtimes_{j \in t(T)} U^{(j)}) \\ \cong &\bigotimes_{A \in \pi_0(T^0)} E^* \bigotimes_{i \rightharpoondown j} \mathcal{E}(U^\star_{(j)}, U_{(i)})^* \bigotimes_{l \in t(T^1)} \mathcal{E}(U_{(\sigma(l))}, U^{(l)}) \bigotimes_{i \rightharpoonup j} \mathcal{E}^{\boxtimes 2}(H, U^{(j)} \boxtimes U^{(i)}). \end{aligned} \tag{45}$$

3.3 DW invariants of $\langle 2\rangle$-3-manifolds

For an ordinary 3-manifold C, re-define $Z(C)$ as follows.

If C is closed, let

$$Z(C) = \frac{1}{\#\Gamma} \cdot \#\pi_0(\mathfrak{B}^\Gamma(C)), \tag{46}$$

which coincides with (2).

If $\partial C = B \neq \emptyset$, for $[\rho] \in \pi_0(\mathfrak{B}^\Gamma(B))$, choose a representative $\rho \in [\rho]$ and put

$$Z(C)([\rho]) = \#\pi_0(\mathfrak{B}^\Gamma_\rho(C)), \tag{47}$$

where $\mathfrak{B}^\Gamma_\rho(C)$ is the subcategory of $\mathfrak{B}^\Gamma(C)$ having objects $\phi : \Pi_1(C) \to \Gamma$ such that $\phi|_B = \rho$ and morphisms those taking e on B. It is easy to see that this is independent of the choice of ρ, and actually coincides with (3) when (25) is identified with (1).

Now turn to more general 3-manifolds.

Definition 3.8. *An n-manifold with corners X is an n-dimensional topological manifold together with an n-dimensional smooth structure with corners which is a maximal family of charts $\{(U_i, \phi_i) \colon i \in I\}$ such that: (i) $\{U_i \colon i \in I\}$ is an open cover of X; (ii) each $\phi_i : U_i \to \mathbb{R}^n_+ := [0, \infty)^n$ is a homeomorphism from U onto an open subset of $\mathbb{R}^n_+$; (iii) for all $i, j \in I$, $\phi_j \phi_i^{-1} : \phi_i(U_i \cap U_j) \to \phi_j(U_i \cap U_j)$ can be extended to a diffeomorphism between open subsets of $\mathbb{R}^n$.*

Remark 3.9. *For manifolds with corners, the meanings of "boundary", "orientation" and "diffeomorphsim" are all self-evident.*

Definition 3.10. *A $\langle 2 \rangle$-*3-manifold *C is a 3-manifold with corners together with cobordisms*

$$B_{h,1} : A_{d,1} \to A_{d,2}, \qquad B_{h,2} : A_{u,1} \to A_{u,2},$$
$$B_{v,1} : A_{d,1} \to A_{u,1}, \qquad B_{v,2} : A_{d,2} \to A_{u,2},$$

such that $\partial C = -B_{h,1} \cup B_{h,2} \cup -B_{v,1} \cup B_{v,2}$. Write C as

$$\begin{tikzcd}
A_{u,1} \arrow[r, "B_{h,2}"] & A_{u,2} \\
A_{d,1} \arrow[u, "B_{v,1}"] \arrow[r, "B_{h,1}"] & A_{d,2} \arrow[u, "B_{v,2}"'] \arrow[ul, Rightarrow, "C"']
\end{tikzcd} \tag{48}$$

Remark 3.11. *We allow some of the involved submanifolds to be empty. When the 1-submanifolds $A_{d,1}, A_{d,2}, A_{u,1}, A_{u,2}$ are all empty, C is just an ordinary 3-manifold probably with boundary.*

Definition 3.12. *A diffeomorphism $F : C \to C'$ of $\langle 2 \rangle$-3-manifolds is a diffeomorphism of manifolds with corners such that $F(B_{v,i}) = B'_{v,i}$, $F(B_{h,i}) = B'_{h,i}$ for $i = 1, 2$.*

For a $\langle 2 \rangle$-3-manifold C as in (48), we define a natural transformation

$$Z(C) : \hat{Z}(B_{v,2}) \circ \hat{Z}(B_{h,1}) \to \hat{Z}(B_{h,2}) \circ \hat{Z}(B_{v,1}),$$

and call the following square the *DW invariant* of C:

$$\begin{tikzcd}
\mathcal{E}^{\boxtimes\pi_0(A_{u,1})} \arrow[r, "Z(B_{h,2})"] & \mathcal{E}^{\boxtimes\pi_0(A_{u,2})} \\
\mathcal{E}^{\boxtimes\pi_0(A_{d,1})} \arrow[u, "Z(B_{v,1})"] \arrow[r, "Z(B_{h,1})"] & \mathcal{E}^{\boxtimes\pi_0(A_{d,2})} \arrow[u, "Z(B_{v,2})"'] \arrow[ul, Rightarrow, "Z(C)"']
\end{tikzcd} \tag{49}$$

The natural transformation $Z(C)$ is induced by a map

$$Z(C) : Z(B_{v,2} \cup B_{h,1}) \to Z(B_{h,2} \cup B_{v,1}) \tag{50}$$

which is a morphism in $\mathcal{E}^{\boxtimes(\pi_0(A_{d,1})\sqcup\pi_0(A_{u,2}))}$, described explicitly as follows.

Suppose

$$\#\pi_0(A_{d,1}) = p, \quad \#\pi_0(A_{d,2}) = q, \quad \#\pi_0(A_{u,1}) = q', \quad \#\pi_0(A_{u,2}) = r.$$

For $\mathbf{x} \in \Gamma^p, \mathbf{y} \in \Gamma^q, \mathbf{z} \in \Gamma^r$, and $[\psi] \in \pi_0(\mathcal{B}^\Gamma_{\mathbf{z},\mathbf{y^{-1}}}(B_{v,2})), [\phi] \in \pi_0(\mathcal{B}^\Gamma_{\mathbf{y},\mathbf{x^{-1}}}(B_{h,1}))$, let $\delta([\phi]) \in Z(B_{h,1})$ take 1 at $[\phi]$ and 0 otherwise; define $\delta([\psi]) \in Z(B_{v,2})$ similarly. Let

$$\delta([\psi],[\phi]) = \sum_{\mathbf{g}\in\Gamma^q} \delta(\eta_{\mathbf{e},\mathbf{g}} \cdot [\psi]) \otimes \delta(\eta_{\mathbf{g},\mathbf{e}} \cdot [\phi]) \in Z(B_{v,2} \cup B_{h,1}). \tag{51}$$

We have

$$\delta([\psi],[\phi]) = \delta(\eta_{\mathbf{e},\mathbf{h}} \cdot [\psi], \eta_{\mathbf{h},\mathbf{e}} \cdot [\phi]), \tag{52}$$

for all $\mathbf{h} \in \Gamma^q$. When $\mathbf{x}, \mathbf{y}, \mathbf{z}$ and $[\phi], [\psi]$ vary, the different element $\delta([\psi],[\phi])$ form a canonical orthonomal basis of $Z(B_{h,2} \cup B_{v,1})$.

Similarly we define $\delta([\psi'],[\phi'])$ for $\mathbf{y'} \in \Gamma^{q'}$ and $[\psi'] \in \pi_0(\mathfrak{B}^\Gamma_{\mathbf{z},\mathbf{y'^{-1}}}(B_{h,2})), [\phi'] \in \pi_0(\mathfrak{B}^\Gamma_{\mathbf{y'},\mathbf{x^{-1}}}(B_{v,1}))$. The various $\delta([\psi'],[\phi'])$ form a canonical orthonomal basis of $Z(B_{h,2} \cup B_{v,1})$.

Now choose representatives $\phi \in [\phi], \psi \in [\psi], \phi' \in [\phi'], \psi' \in [\psi']$, and let $\mathfrak{B}^{\psi',\phi'}_{\psi,\phi}(C)$ be the subcategory of $\mathfrak{B}^\Gamma(C)$ having objects $\rho : \Pi_1(C) \to \Gamma$ such that $\rho|_{\Pi_1(B_{h,1})} = \phi, \rho|_{\Pi_1(B_{v,2})} = \psi, \rho|_{\Pi_1(B_{v,1})} = \phi', \rho|_{\Pi_1(B_{h,2})} = \psi'$ and morphisms $\eta : \rho \to \rho'$ satisfying $\eta|_{\partial C} \equiv e$. Set

$$Y(C)^{[\psi'],[\phi']}_{[\psi],[\phi]} = \#\pi_0(\mathfrak{B}^{\psi',\phi'}_{\psi,\phi}(C)); \tag{53}$$

it depends only on the classes $[\psi], [\phi], [\psi'], [\phi']$ hence is well-defined. Define

$$Z(C)(\delta([\psi],[\phi])) = \sum Y(C)^{[\psi'],[\phi']}_{[\psi],[\phi]} \cdot \delta([\psi'],[\phi']), \tag{54}$$

the sum taking over the canonical basis of $Z(B_{v,2} \cup B_{h,1})$. It is easy to verify that $Z(C)$ is indeed a morphism in $\mathcal{E}^{\boxtimes(p+r)}$.

Theorem 3.13. *(a) (Functoriality) For each orientation-preserving diffeomorphism of $\langle 2\rangle$-3-manifolds $F : C \to C'$, the following diagram commutes*

$$\begin{tikzcd}
\hat{Z}(B_{v,2}) \circ \hat{Z}(B_{h,1}) \arrow[r, "Z(C)"] \arrow[d, "(\partial F)_*"'] & \hat{Z}(B_{h,2}) \circ \hat{Z}(B_{v,1}) \arrow[d, "(\partial F)_*"] \\
\hat{Z}(B'_{v,2}) \circ \hat{Z}(B'_{h,1}) \arrow[r, "Z(C')"] & \hat{Z}(B'_{h,2}) \circ \hat{Z}(B'_{v,1})
\end{tikzcd}$$

(b) (Multiplicativity) For two $\langle 2\rangle$-3-manifolds C and C', one has $Z(C \sqcup C') = Z(C) \boxtimes Z(C')$.

(c) (Gluing) Given two $\langle 2\rangle$-3-manifolds

$$\begin{tikzcd}
A_{u,1} \arrow[r, "B_{h,2}"] & A_{u,2} & & A'_{u,1} \arrow[r, "B'_{h,2}"] & A'_{u,2} \\
A_{d,1} \arrow[u, "B_{v,1}"] \arrow[r, "B_{h,1}"] & A_{d,2} \arrow[u, "B_{v,2}"'] \arrow[ul, Rightarrow, "C"'] & & A'_{d,1} \arrow[u, "B'_{v,1}"] \arrow[r, "B'_{h,1}"] & A'_{d,2} \arrow[u, "B'_{v,2}"'] \arrow[ul, Rightarrow, "C'"']
\end{tikzcd}$$

(i) If $B_{h,2} = B'_{h,1}$ (so that $A_{u,1} = A'_{d,1}, A_{u,2} = A'_{d,1}$), one can glue C and C' along $B_{h,2}$ to get a new $\langle 2\rangle$-3-manifold C_v, then for all $U \in Z(A_{d,1})$,

$$Z(C_v)_U = Z(C')_{\hat{Z}(B_{v,1})(U)} \circ \hat{Z}(B'_{v,2})(Z(C)_U) :$$
$$(\hat{Z}(B'_{v,2} \cup B_{v,2}) \circ \hat{Z}(B_{h,1}))_U \to (\hat{Z}(B'_{h,2}) \circ \hat{Z}(B'_{v,1} \cup B_{v,1}))_U.$$

(ii) If $B_{v,2} = B'_{v,1}$ (so that $A_{u,2} = A'_{u,1}, A_{d,2} = A'_{d,1}$), one can glue C and C' along $B_{v,2}$ to get a new $\langle 2\rangle$-3-manifold C_h, then for all $U \in Z(A_{d,1})$,

$$Z(C_h)_U = \hat{Z}(B'_{h,2})(Z(C)_U) \circ Z(C')_{\hat{Z}(B_{h,1})(U)} :$$
$$(\hat{Z}(B'_{v,2}) \circ \hat{Z}(B'_{h,1} \cup B_{h,1}))_U \to (\hat{Z}(B'_{h,2} \cup B_{h,2}) \circ \hat{Z}(B_{v,1}))_U$$

From now on in this section, we assume that all connected $\langle 2\rangle$-3-manifolds C satisfy $B_{v,2} = CL$, so $Z(B_{v,2}) \cong \mathrm{id}_{\mathcal{E}}$.

Suppose C is connected. Given $U \in Z(A_{d,1}), U' \in Z(A_{u,1})$, and $\alpha : \hat{Z}(B_{v,1})(U) \to U'$, there is an induced map

$$Z(C)_\alpha := \hat{Z}(B_{h,2})(\alpha) \circ Z(C)_U : \hat{Z}(B_{h,1})(U) \to \hat{Z}(B_{h,2})(U'). \tag{55}$$

Proposition 3.14. *(a) Suppose C, C' are connected $\langle 2\rangle$-3-manifolds, and $F : C \to C'$ is an orientation-preserving diffeomorphism. Given $U \in Z(A_{d,1}), U' \in Z(A_{u,1})$, and $\alpha : \hat{Z}(B_{v,1})(U) \to U'$, let α' be the composite*

$$\hat{Z}(B'_{v,1})(U) \xrightarrow{((F^{-1}|_{B'_{v,1}})_*)_U} \hat{Z}(B_{v,1})(U) \xrightarrow{\alpha} U',$$

then we have the following commuting diagram

$$\begin{tikzcd}
\hat{Z}(B_{h,2})(U') \arrow[r, "((F|_{B_{h,2}})_*)_{U'}"] & \hat{Z}(B'_{h,2})(U') \\
\hat{Z}(B_{h,1})(U) \arrow[r, "((F|_{B_{h,1}})_*)_{U}"] \arrow[u, "Z(C)_\alpha"] & \hat{Z}(B'_{h,1})(U) \arrow[u, "Z(C')_{\alpha'}"']
\end{tikzcd}$$

(b) Suppose C, C' are connected $\langle 2\rangle$-3-manifolds satisfying the conditions of Proposition 3.13(c)(i). For any

$$U \in Z(A_{d,1}), \quad U' \in Z(A_{u,1}), \quad U'' \in Z(A'_{u,1}),$$

$$\alpha : \hat{Z}(B_{v,1})(U) \to U', \quad \alpha' : \hat{Z}(B'_{v,1})(U') \to U'',$$

we have

$$Z(C_v)_{\alpha' \circ \hat{Z}(B'_{v,1})(\alpha)} = Z(C')_{\alpha'} \circ Z(C)_\alpha.$$

(c) Suppose C^1, C^2 are connected $\langle 2\rangle$-3-manifolds. Glue $C^1 \sqcup C^2$ with $C(\mathbf{0})$ by identifying $B^1_{v,2} \sqcup B^2_{v,2}$ with the $B_{v,1}(\mathbf{0})$, to get a new $\langle 2\rangle$-3-manifold $C^1 \odot C^2$. Then for all

$$U_i \in Z(A^i_{d,1}), \quad U'_i \in Z(A^i_{u,1}), \quad \alpha^i : \hat{Z}(B^i_{v,1})(U_i) \to U'_i, \quad i = 1, 2,$$

we have

$$Z(C^1 \odot C^2)_{\alpha^1 \boxtimes \alpha^2} = Z(C^1)_{\alpha^1} \odot Z(C^2)_{\alpha^2}.$$

3.4 Examples II



Figure 7: (a) $C(\mathbf{0})$; (b) $C(\cap)$; (c) $C(\mathbf{1})$

Let $C(\mathbf{0}), C(\cap), C(\mathbf{1})$ be the $\langle 2\rangle$-3-manifolds shown in Figure 7.

It is obvious that $Z(C(\mathbf{0}))$ is the identity natural transformation.

The $\langle 2\rangle$-3-manifolds $C(\cap), C(\mathbf{1})$ are presented as squares as follows:

$$\begin{tikzcd}
\emptyset \arrow[r, "D"] & S^1 \\
2S^1 \arrow[u, "B(\cap)"] \arrow[r, "P"] & S^1 \arrow[u, "CL"'] \arrow[ul, Rightarrow, "C(\cap)"']
\end{tikzcd}
\qquad
\begin{tikzcd}
2S^1 \arrow[r, "P"] & S^1 \\
2S^1 \arrow[u, "B(\mathbf{1})"] \arrow[r, "P"] & S^1 \arrow[u, "CL"'] \arrow[ul, Rightarrow, "C(\mathbf{1})"']
\end{tikzcd}$$

(1) By Section 3.2,

$$\begin{aligned}
\hat{Z}(CL)\hat{Z}(P)(U\boxtimes V)_z &\cong \frac{1}{(\#\Gamma)^2}\Big(\bigoplus_{\substack{a_1,a_2,x_1,x_2:\\ {}^{a_1}x_1\cdot{}^{a_2}x_2=z}} \mathbb{C}\langle x_1^{-1},x_2^{-1},a_1,a_2\rangle\otimes U_{x_1}\otimes V_{x_2}\Big)^{\Gamma^2} \\
&\cong (U\odot V)_z && (56)\\
\hat{Z}(B_{h,2})\hat{Z}(B_{v,1})(U\boxtimes V)_z &\cong \frac{1}{(\#\Gamma)^2}\Big(\bigoplus_{\substack{b_1,b_2,x_1,x_2:\\ {}^{b_1}x_2\cdot{}^{b_2}x_1=z}} \mathbb{C}\langle x_2^{-1},x_1^{-1},b_1,b_2\rangle\otimes U_{x_1}\otimes V_{x_2}\Big)^{\Gamma^2} \\
&\cong (V\odot U)_z, && (57)
\end{aligned}$$

where we have set the holonomies along non-horizontal paths to be identity, hence $y_1=x_2, y_2=x_1$. From (35) we see that, the image of $u_{x_1}\otimes v_{x_2}\in(U\otimes V)_z$ under the inverse of (56) is

$$\sum_{a_1,a_2}\langle {}^{a_1^{-1}}x_1^{-1}, {}^{a_2^{-1}}x_2^{-1}, a_1, a_2\rangle\otimes(a_1^{-1}u_{x_1})\otimes(a_2^{-1}v_{x_2}),$$

which is then sent by $Z(C)_{U\boxtimes V}$ to

$$\sum_{a_1,a_2}\langle {}^{a_2^{-1}}x_2^{-1}, {}^{a_1^{-1}}x_1^{-1}, {}^{a_1}x_1a_2, a_1\rangle\otimes(a_2^{-1}v_{x_2})\otimes(a_1^{-1}u_{x_1}), \qquad (58)$$

since it is forced that $b_2=a_1, b_1={}^{a_1}x_1a_2$; finally, by (35), the map (57) sends (58) to $(x_1v_{x_2})\otimes u_{x_1}$. Thus

$$Z(C(\mathbf{1}))_{U\boxtimes V}=R_{U,V}. \qquad (59)$$

(2)

$$\begin{aligned}
\hat{Z}(CL)\hat{Z}(P)(U\boxtimes V) &\cong U\odot V, && (60)\\
\hat{Z}(D)\hat{Z}(B(\cap))(U\boxtimes V) &\cong \mathcal{E}(U^\star, V). && (61)
\end{aligned}$$

The image of $u\otimes v$ under the inverse of (60) is

$$\sum_{a_1,a_2}\langle {}^{a_1^{-1}}x_1^{-1}, {}^{a_2^{-1}}x_2^{-1}, a_1, a_2\rangle\otimes a_1^{-1}u\otimes a_2^{-1}v,$$

and then is sent by $Z(C(\cap))_{U\boxtimes V}$ to

$$\sum_{a_1,a_2}\langle a_2^{-1}x_2^{-1}, a_1^{-1}a_2\rangle\otimes a_1^{-1}u\otimes a_2^{-1}v,$$

using the relation $g = a_1^{-1}a_2$; finally, by (35) and (38), it is sent by (61) to the morphism

$$\overline{u'} \mapsto \sum_g (gu, u')_U \cdot gv.$$

Hence

$$Z(C(\cap))_{U \boxtimes V} : U \odot V \to \mathcal{E}(V^\star, U), \tag{62}$$

$$u \otimes v \mapsto (\overline{u'} \mapsto \sum_g (gu, u')_U \cdot gv). \tag{63}$$

(3) Let $C(\cup)$ be the $\langle 2 \rangle$-3-manifold obtained by reflecting $C(\cap)$ along the horizontal plane. Then by a similar manner, we obtain

$$Z(C(\cup))_{\mathbb{C}} : \underline{\mathbb{C}} \to \hat{Z}(P)(H) = \bigoplus_z (\bigoplus_{xy=z} H(y, x), \tag{64}$$

$$1 \mapsto \sum_x \langle x, e \rangle. \tag{65}$$

(4) Suppose L is a framed link, and let $C = C(L)$ denote the complement of a tubular neighborhood $N(L)$ of L in a solid cylinder containing it; this is a $\langle 2 \rangle$-3-manifold with $B_{h,1} = B_{h,2} = D$, $B_{v,1} = L \times S^1, B_{v,2} = CL$.

By definition, $Z(C(L))_{\mathbb{C}} : \underline{\mathbb{C}} \to \underline{Z(\partial N(L))}$ is the map sending 1 to

$$\sum_{[\rho] \in \pi_0(\mathfrak{B}^\Gamma(\partial N(L)))} \# \pi_0(\mathfrak{B}^\Gamma_{\rho,e}(C(L))) \cdot [\rho],$$

where ρ is an arbitrary representative of $[\rho]$, $\mathfrak{B}^\Gamma_{\rho,e}(C(L))$ is the subcategory of $\mathfrak{B}^\Gamma(C(L))$ having objects $\Phi : \Pi_1(C(L)) \to \Gamma$ such that $\Phi|_{\Pi_1(\partial C(L) - \partial N(L))} = e, \Phi|_{\Pi_1(\partial N(L))} = \rho$ and morphisms $\eta : C(L) \to \Gamma$ taking e on $\partial C(L)$.

Note that $\partial C(L)$ is homeomorphic to S^2. Identifying $W \in \mathcal{V}$ with $\underline{W}$, then

$$Z(C(L))_{\mathbb{C}}(1) \otimes 1 = Z(D^3 - N(L)) \in Z(\partial N(L)) \otimes Z(-S^2).$$

Hence, using (4) and applying the gluing axiom (D) of Section 2.1, we have

$$Z(S^3 - N(L)) = (Z(D^3 - N(L)), Z(D^3))_{Z(S^2)} = \frac{1}{\#\Gamma} Z(C(L))_{\mathbb{C}}(1). \tag{66}$$

(5) Define

$$\epsilon_U : \hat{Z}(B(\cap))(U^\star \boxtimes U) \cong \mathrm{End}_{\mathcal{E}}(U) \to \mathbb{C}, \quad \mathrm{id}_U \mapsto \frac{\mathrm{Dim} U}{\#\Gamma}; \tag{67}$$

$$\iota_U : \hat{Z}(B(\cup))(\mathbb{C}) = H \to U \boxtimes U^\star, \quad \langle x, g \rangle \mapsto \sum_i u_x^i \otimes \overline{g u_x^i}, \tag{68}$$

where $\{u_x^i\}$ is an arbitrary orthonormal basis of U_x.

By definition, $Z(C(\cap))_{\epsilon_U}$ is the composite

$$U^\star \odot U \xrightarrow{Z(C(\cap))_{U^\star \boxtimes U}} \underline{\mathrm{End}_{\mathcal{E}}(U)} \xrightarrow{\hat{Z}(D)(\epsilon_U)} \underline{\mathbb{C}},$$

and $Z(C(\cup))_{\iota_U}$ is the composite

$$\underline{\mathbb{C}} \xrightarrow{Z(C(\cup))_{\mathbb{C}}} \hat{Z}(P)(H) \xrightarrow{\hat{Z}(P)(\iota_U)} U \odot U^\star.$$

It can be verified that

$$Z(C(\cap))_{\epsilon_U} = e_U, \qquad Z(C(\cup))_{\iota_U} = i_U. \tag{69}$$

(6) Consider the $\langle 2\rangle$-3-manifold $C(\mathbf{1})$. we have

$$Z(C(\mathbf{1}))_{\mathrm{id}_{V\boxtimes U}} = R_{U,V}. \tag{70}$$

4 Relating DW invariants to RT invariants

From each framed tangle T, one can construct a connected $\langle 2\rangle$-3-manifold by removing a tubular neighborhood of T from a sufficiently large solid cylinder containing T; denote it as

$$\begin{tikzcd}
A_{u,1}(T) \arrow[r, "B_{h,2}(T)"] & A_{u,2}(T) \\
A_{d,1}(T) \arrow[u, "B_{v,1}(T)"] \arrow[r, "B_{h,1}(T)"'] & A_{d,2}(T) \arrow[u, "B_{v,2}(T)"'] \arrow[ul, Rightarrow, "C(T)"']
\end{tikzcd} ;$$

obviously $B_{v,2}(T) = CL$.

Given a colored framed tangle (T,c), by Definition 2.3 we can always assume that for all $A \in T^\pm$, the terminal point is on the left of the initial point. For $A \in \pi_0(T)$, define $\tilde{\alpha}_{A,c}$ as follows.

- If $A \in \pi_0(T^0)$, let $\tilde{\alpha}_{A,c} \in E^*$ sends χ to $(\chi, \chi_{c(A)})_E$;
- if $A \in \pi_0(T^-)$, put $\tilde{\alpha}_{A,c} = \epsilon_{c(A)} \in \mathrm{End}_{\mathcal{E}}(c(A))^*$;
- if $A \in \pi_0(T^+)$, put $\tilde{\alpha}_{A,c} = \iota_{c(A)}$;
- if $A \in \pi_0(T^1)$, put $\tilde{\alpha}_{A,c} = \mathrm{id}_{c(A)}$.

Suppose $\#s(T) = k, \#t(T) = l$, let $U_{(i)}, i = 1, \cdots, k; U^{(j)}, j = 1, \cdots, l$ and U_T, U^T be the objects defined in Section 2.4. Let $\alpha_{T,c}$ be the inverse image of the morphism

$$\tilde{\alpha}_{T,c} = \boxtimes_{A\in\pi_0(T)} \tilde{\alpha}_{A,c} \tag{71}$$

under (45). Note that $\hat{Z}(B_{h,1})(U_{(1)} \boxtimes \cdots \boxtimes U_{(k)}) = U_T$ and $\hat{Z}(B_{h,2})(U^{(1)} \boxtimes \cdots \boxtimes U^{(l)}) = U^T$. Define

$$F(T,c) = Z(C(T))_{\alpha_{T,c}} : U_T \to U^T. \tag{72}$$

By Proposition 3.14 (a), this is an invariant of (T,c).

Lemma 4.1. *Given two colored framed tangles* (T_1,c_1) *and* (T_2,c_2)*, we have*

$$\alpha_{T_1 \sqcup T_2, c_1 \sqcup c_2} = \alpha_{T_1,c_1} \boxtimes \alpha_{T_2,c_2}; \tag{73}$$

if $t(T_1,c_1) = s(T_2,c_2)$*, then*

$$\alpha_{T_2 \cdot T_1, c_2 \cdot c_1} = \alpha_{T_2,c_2} \circ \hat{Z}(B_{v,1}(T_2))(\alpha_{T_1,c_1}). \tag{74}$$

Proof. The first equation $\alpha_{T_1 \sqcup T_2, c_1 \sqcup c_2} = \alpha_{T_1,c_1} \boxtimes \alpha_{T_2,c_2}$ is obvious.

For the second, it suffices to show

$$\tilde{\alpha}_{T_2 \cdot T_1, c_2 \cdot c_1} = \tilde{\alpha}_{T_2,c_2} \circ (\hat{Z}^0(T_2) \otimes \hat{Z}'(T_2))(\tilde{\alpha}_{T_1,c_1}). \tag{75}$$

Without loss of generality, we can assume $\#t(T_1) = \#s(T_2) = 2$ and one of the following cases occurs: 1. $T_1 = T_1^+, T_2 = T_2^1$; 2. $T_1 = T_1^+, T_2 = T_2^-$; 3. $T_1 = T_1^1, T_2 = T_2^-$; 4. $T_1 = T_1^1, T_2 = T_2^1$.

The cases 3 and 4 are trivial. For 1, just note that $\iota_{c_1(T_1)}$ under (41) corresponds to $\mathrm{id}_{c_1(T_1)}$. To verify (75) for case 2, suppose $c_1(T_1) = U$, then

$$c_2(T_2) = U^\star, \qquad \tilde{\alpha}_{T_1,c_1} = \iota_U, \qquad \tilde{\alpha}_{T_2,c_2} = \epsilon_{U^\star}.$$

Note that, by (42) and (38),

$$\hat{Z}'(T_2)(\iota_U) : E \cong \hat{Z}'(T_2)(H) \to \mathrm{End}_{\mathcal{E}}(U^\star)$$

sends $\chi = \sum_{x,g} \chi(x,g) \cdot \langle x,g \rangle$ to $(\overline{u} \mapsto \sum_{x,g} \chi(x,g) \sum_i (u_x^i, u)_U \cdot \overline{g u_x^i})$, which is actually

$$\Big(\frac{1}{\mathrm{Dim} U} \sum_{x,g} \chi(x,g) \overline{\chi_U(x,g)}\Big) \cdot \mathrm{id}_{U^\star}, \tag{76}$$

obtained by computing its trace; then (76) is sent by $\epsilon_{U^\star}$ to $(\chi, \chi_U)_E$, which is equal to $\tilde{\alpha}_{T_2 \cdot T_1, c_2 \cdot c_1}(\chi)$. □

This lemma together with Proposition 3.14 (b), (c) imply

Proposition 4.2. *For two colored framed tangles* (T_1,c_1) *and* (T_2,c_2)*.*

(a) we have

$$F(T_1 \sqcup T_2, c_1 \sqcup c_2) = F(T_1,c_1) \odot F(T_2,c_2).$$

(b) If $t(T_1,c_1) = s(T_2,c_2)$*, then*

$$F(T_2 \cdot T_1, c_2 \cdot c_1) = F(T_2,c_2) \circ F(T_1,c_1).$$

Since by Example (5), (6) of Section 3.4, F and $\mathcal{F}$ associate the same morphism to each colored basic tangle, we have

Corollary 4.3. *For all colored framed tangles* (T, c), *we have*

$$F(T, c) = \mathcal{F}(T, c).$$

Theorem 4.4. *For all framed link* $L = K_1 \sqcup \cdots \sqcup K_r$,

$$Z(L) = \frac{1}{\#\Gamma} \sum_{U(1),\cdots,U(r)\in\Lambda} F(L, c(U(1), \cdots, U(r))) \cdot \bigotimes_{i=1}^{r} \chi_{U(i)}, \tag{77}$$

where the coloring $c(U(1), \cdots, U(r))$ *takes* $U(i)$ *at* K_i *for* $1 \leqslant i \leqslant r$.

Proof. Abbreviate $c(U(1), \cdots, U(r))$ to c. By definition, $F(L, c) : \mathbb{C} \to \mathbb{C}$ is the composite of the upper row of the following commuting diagram

$$\begin{tikzcd}
\mathbb{C} \arrow[r, "Z(C(L))_{\mathbb{C}}"] \arrow[d, "="] & Z(L \times S^1) \arrow[r, "\alpha_{L,c}"] \arrow[d, "\cong"] & \mathbb{C} \arrow[d, "="] \\
\mathbb{C} \arrow[r] & E^{\otimes r} \arrow[r, "\tilde{\alpha}_{L,c}"] & \mathbb{C}
\end{tikzcd} \tag{78}$$

where the middle isomorphism is induced by the homeomorphism $L \times S^1 \cong \sqcup_r \Sigma_1$ determined by the framing of L. Now $Z(C(L))_{\mathbb{C}}(1) = \#\Gamma \cdot Z(S^3 - N(L))$ by (66), it is then sent by the middle isomorphism to $\#\Gamma \cdot Z(L)$, and finally sent by $\tilde{\alpha}_{L,c}$ to $\#\Gamma \cdot (Z(L), \chi_c)_{E^{\otimes r}}$, with $\chi_c = \bigotimes_{i=1}^{r} \chi_{U(i)}$.

The result follows. ☐

5 Computing invariants of Montesinos knots

For a framed tangle T, let $r(T)$ denote the new framed tangle represented by the tangle diagram obtained by rotating that of T counterclockwise by $\frac{\pi}{2}$.

For integers $s_1, \cdots, s_k$, define the *continued fraction* $[[s_1, \cdots, s_k]]$ by

$$[[s_1]] = s_1, \qquad [[s_1, \cdots, s_{r-1}, s_k]] = s_k - 1/[[s_1, \cdots, s_{k-1}]]. \tag{79}$$

For a rational number $\frac{p}{q} \in \mathbb{Q}$ (with $(p, q) = 1$), the *rational tangle* $T(\frac{p}{q})$ is $r(\mathbf{c_k} \cdot (\cdots r(\mathbf{c_2} \cdot r(\mathbf{c_1})) \cdots))$, where the c_i are chosen so that $[[c_1, \cdots, c_k]] = \frac{p}{q}$.

Remark 5.1. *It can be proved by induction on* k *that any rational tangle is invariant under the rotation by* π *along each of the axes of* $\mathbb{R}^3$. *Thus* $r(T(\frac{p}{q})) = r'(T(\frac{p}{q}))$ *where* r' *is the clockwise rotation by* $\frac{\pi}{2}$, *and our definition of rational tangle coincides with that given on Wikipedia.*

A *Montesinos tangle* is the composite tangle $T(\frac{p_m}{q_m})\cdots T(\frac{p_1}{q_1})$ for some rational numbers $\frac{p_1}{q_1},\cdots,\frac{p_m}{q_m}$, and its closure is called a *Montesinos link* and denoted as $M(\frac{p_1}{q_1},\cdots,\frac{p_m}{q_m})$. In particular, $M(\frac{p}{q})$ is called a *rational link* (another name is *2-bridge link*), and $M(p_1,\cdots,p_m)$ is called a *pretzel link*. A $\mathcal{P}$ knot is a $\mathcal{P}$ link with one component, for arbitrary adjective $\mathcal{P}$.

In this section we shall see that the problem of computing the DW invariants of Montesinos knots can be reduced to a linear-algebraic one.

Suppose (T,c) is a colored framed tangle such that $\#s(T)=\#t(T)=2$ as shown on the left of Figure 8; associated to it is a morphism $F(T,c): U_L \odot U_R \to U^L \odot U^R$.



Figure 8: The reflection

From the figure we see that $F(r(T,c)) : (U^L)^\star \odot U_R \to U^R \odot U_R^\star$ is equal to the composite

$$(e_U \odot \mathrm{id}_{U^R} \odot \mathrm{id}_{U_R^\star}) \circ (\mathrm{id}_{(U^L)^\star} \odot F(T,c) \odot \mathrm{id}_{U_R^\star}) \circ (\mathrm{id}_{(U^L)^\star} \odot \mathrm{id}_{U_L} \odot i_{U_R}).$$

Define

$$\mathrm{rot} = \mathrm{rot}_{U_1,U_2}^{U_3,U_4} : \mathcal{E}(U_1 \odot U_2, U_3 \odot U_4) \to \mathcal{E}(U_3^\star \odot U_1, U_4 \odot U_2^\star), \tag{80}$$

$$F \mapsto (e_{U_3} \odot \mathrm{id}_{U_4} \odot \mathrm{id}_{U_2^\star}) \circ (\mathrm{id}_{(U_3)^\star} \odot F \odot \mathrm{id}_{U_2^\star}) \circ (\mathrm{id}_{U_3^\star} \odot \mathrm{id}_{U_1} \odot i_{U_2}). \tag{81}$$

Then

$$F(r(T,c)) = \mathrm{rot}(F(T,c)). \tag{82}$$

Definition 5.2. *For $W \in \mathcal{E}$ and $f \in End_{\mathcal{E}}(W)$, the* vector-valued trace $\overrightarrow{tr}(f)$ *is the element of E given by,*

$$\overrightarrow{tr}(f) = \sum_{V \in \Lambda} tr(f_* : \mathcal{E}(V,W) \to \mathcal{E}(V,W)) \cdot \chi_V. \tag{83}$$

For $w \in E$, define

$$|w| = \sum_{V \in \Lambda} DimV(w, \chi_V)_E. \tag{84}$$

Lemma 5.3. *If the colored framed tangle (T, c) satisfies $s(T) = t(T)$ and $U_T = U^T$, we can get a colored framed link $(\overline{T}, \overline{c})$ by taking the closure. Then*

$$F(\overline{T}, \overline{c}) = |\overrightarrow{tr}(F(T, c))|. \tag{85}$$

Proof. By Lemma 2.6 of [20], $\mathcal{F}(\overline{T}, \overline{c}) = \mathrm{tr}_q \mathcal{F}(T, c)$, where tr_q, in our setting and notation, is equal to $|\overrightarrow{tr}(F(T, c))|$, as can be checked. □

Theorem 5.4. *(1) Suppose $\frac{p}{q} = [c_1, \cdots, c_k]$, then*

$$F(T(\frac{p}{q}), U) = rot(R^{c_k} \cdot (\cdots rot(R^{c_1}))). \tag{86}$$

(2) The DW invariant of the Montesinos knot $K = M(\frac{p_1}{q_1}, \cdots, \frac{p_m}{q_m})$ can be expressed as

$$Z(K) = \frac{1}{\#\Gamma} \sum_{U \in \Lambda} F(K, U) \cdot \chi_U, \tag{87}$$

$$F(K, U) = \big|\overrightarrow{tr}(F(T(\frac{p_m}{q_m}), U) \circ \cdots \circ F(T(\frac{p_1}{q_1}), U))\big|. \tag{88}$$

Remark 5.5. *By (T, U) we mean the colored framed tangle with the coloring taking U at all components. By R we mean $R_{V,W}$ for some $V, W \in \{U, U^\star\}$ as determined by the colored framed tangle; it should be warned that the many R may differ from each other. Similarly for rot.*

We now show how to compute $Z(M(\frac{p_1}{q_1}, \cdots, \frac{p_m}{q_m}))$ practically.

Step 1. For all $U, V \in \Lambda$, choose isomorphisms

$$\beta_{U,V} : U \odot V \cong \bigoplus_{W \in \Lambda} N_{U,V}^W \cdot W, \tag{89}$$

with $N_{U,V}^W \cdot W$ meaning the direct sum of $N_{U,V}^W$ copies of W.

For $U_1, U_2, U_3, U_4 \in \Lambda$, let $\Phi = \Phi_{U_1,U_2}^{U_3,U_4}$ be the composite

$$\begin{aligned} \mathcal{E}(U_1 \odot U_2, U_3 \odot U_4) &\cong \bigoplus_{W \in \Lambda} \mathcal{E}(N_{U_1,U_2}^W W, N_{U_3,U_4}^W W) && (90) \\ &\cong \bigoplus_{W \in \Lambda} \mathcal{V}(\mathbb{C}^{N_{U_1,U_2}^W}, \mathbb{C}^{N_{U_3,U_4}^W}) && (91) \\ &\cong \bigoplus_{W \in \Lambda} \mathcal{M}(N_{U_3,U_4}^W, N_{U_1,U_2}^W), && (92) \end{aligned}$$

where (90) sends f to $\beta_{U_3,U_4}\circ f\circ\beta_{U_1,U_2}^{-1}$, and in 92 $\mathcal{M}(N,L)$ denotes the space of $N\times L$ matrices over $\mathbb{C}$. We shall abbreviate $\Phi_{U_1,U_2}^{U_3,U_4}$ to Φ whenever there is no danger of confusion.

Step 2. Define $\mathrm{ROT}=\mathrm{ROT}_{U_1,U_2}^{U_3,U_4}$ to be the unique linear map making the following diagram commute:

$$\begin{tikzcd}
\mathcal{E}(U_1\odot U_2,U_3\odot U_4) \arrow[r, "\mathrm{rot}"] \arrow[d, "\Phi"'] & \mathcal{E}(U_3^{\star}\odot U_1,U_4\odot U_2^{\star}) \arrow[d, "\Phi"] \\
\bigoplus\limits_{W\in\Lambda}\mathcal{M}(N_{U_3,U_4}^{W},N_{U_1,U_2}^{W}) \arrow[r, "\mathrm{ROT}"] & \bigoplus\limits_{W\in\Lambda}\mathcal{M}(N_{U_4,U_2^{\star}}^{W},N_{U_3^{\star},U_1}^{W})
\end{tikzcd} \tag{93}$$

Choose orthonormal base $\{u_{1,j}\},\{u_{2,k}\},\{u_{3,j'}\},\{u_{4,k'}\}$ for U_1,U_2,U_3,U_4, respectively. Then (recalling (19)) $\{\overline{u_{2,k}}\}$ is the dual basis for $U_2^{\star}$ and so on.

Suppose

$$F:U_1\odot U_2\to U_3\odot U_4,\qquad u_{1,j}\otimes u_{2,k}\mapsto\sum_{j',k'}\mathcal{T}(F)_{j,k}^{j',k'}\cdot u_{3,j'}\otimes u_{4,k'},\tag{94}$$

then $\mathrm{rot}(F)$ can be expressed as

$$\begin{aligned}\overline{u_{3,i}}\otimes u_{1,j}&\mapsto\sum_{k}\overline{u_{3,i}}\otimes u_{1,j}\otimes u_{2,k}\otimes\overline{u_{2,k}}\\&\mapsto\sum_{k,j',k'}\mathcal{T}(F)_{j,k}^{j',k'}\cdot\overline{u_{3,i}}\otimes u_{3,j'}\otimes u_{4,k'}\otimes\overline{u_{2,k}}\\&\mapsto\sum_{k,k'}\mathcal{T}(F)_{j,k}^{i,k'}\cdot u_{4,k'}\otimes\overline{u_{2,k}}.\end{aligned}$$

Hence with respect to the base $\{\overline{u_{3,i}}\},\{u_{1,j}\},\{u_{4,k}\},\{\overline{u_{2,l}}\}$, $\mathrm{rot}(F)$ is represented by the matrix $\mathcal{T}(\mathrm{rot}(F))$ with

$$\mathcal{T}(\mathrm{rot}(F))_{i,j}^{k,l}=\mathcal{T}(F)_{j,l}^{i,k},\tag{95}$$

and the linear map ROT can be determined according to this relation.

Step 3. Let $\check{F}(T(\frac{p}{q}),U)$ be the map obtained by replacing rot with ROT and R with $\Phi(R)$ in (86), then

$$\begin{aligned}&\overrightarrow{tr}(F(T(\frac{p_m}{q_m}),U)\circ\cdots\circ F(T(\frac{p_1}{q_1}),U))\\=&\overrightarrow{Tr}(\check{F}(T(\frac{p_m}{q_m}),U)\circ\cdots\circ\check{F}(T(\frac{p_1}{q_1}),U)),\end{aligned}\tag{96}$$

where for $G=\bigoplus\limits_{W\in\Lambda}G_W\in\bigoplus\limits_{W\in\Lambda}\mathcal{M}(N^W,N^W)$,

$$\overrightarrow{Tr}(G)=\sum_{W\in\Lambda}\mathrm{tr}(G_W)\cdot\chi_W\in E.\tag{97}$$

Then $Z(M(\frac{p_1}{q_1},\cdots,\frac{p_m}{q_m}))$ can be obtained via (87), (88).

6 The number of n-colorings for n

Let L be an oriented link, presented as a diagram $\tilde{L}$ on the plane; let $\mathrm{Ar}(\tilde{L})$ denote the set of arcs of $\tilde{L}$. Given an odd natural number n, an *n-coloring* of $\tilde{L}$ is a map $\mathrm{Ar}(\tilde{L}) \to \mathbb{Z}/n\mathbb{Z}$ such that, at each crossing, the sum of the values of the under-crossing arcs is equal to twice of the value of the over-crossing arc. It is well-known that there is a bijection between the set of n-colorings of $\tilde{L}$ and the set of homomorphisms $\pi_1(L) \to D_n$ sending the meridian to an element of order two. So the number of n-colorings of $\tilde{L}$ depends only on L itself. Call it the *number of n-colorings* of L and denote it by $\mathrm{CN}_n(L)$.

The number of n-colorings of a link is an invariant with long history. It was initiated by Fox [4], and further studied by Przytycki [19]. There are found to be interesting relations between $\mathrm{CN}_n(L)$ for some n and quantum invariants. For instance, (see [17] Problem 4.16), $\mathrm{CN}_3(L)$ is related to the Jones polynomial via $\mathrm{CN}_3(L) = 3|V_L(e^{\pi\sqrt{-1}/3})|^2$, and a formula relating $\mathrm{CN}_5(L)$ to the Kauffman polynomial is given by $\mathrm{CN}_5(L) = 5|F_L(1, e^{2\pi\sqrt{-1}/5} + e^{-2\pi\sqrt{-1}/5})|^2$. However, despite the elementariness, for general n, there is never seen a formula expressing the number of n-colorings, even for classical pretzel knots.

Now for a knot K, by Remark 2.2,

$$\mathrm{CN}_n(K) = \sum_{k=0}^{n-1}((Q_*^{\mathrm{wr}(K)}Z(K))(a^kb, e) + (Q_*^{\mathrm{wr}(K)}Z(K))(a^kb, a^kb)). \tag{98}$$

6.1 Simple objects of $\mathcal{E}$ for D_n

The dihedral group has a presentation

$$D_n = \langle a, b | a^n, b^2, (ab)^n \rangle.$$

There are $\frac{n+3}{2}$ conjugacy classes:

$$\{e\}; \qquad \{a^{\pm j}\}, j = 1, \cdots, \frac{n-1}{2}; \qquad \{a^kb \colon 0 \leqslant k \leqslant n-1\}.$$

The corresponding centralizers are

$$C(e) = D_n; \qquad C(a^{\pm j}) = \langle a \rangle \cong \mathbb{Z}/n\mathbb{Z}; \qquad C(a^kb) = \langle a^kb \rangle \cong \mathbb{Z}/2\mathbb{Z}.$$

Let X be the trivial representation of D_n, and Y be the 1-dimensional representation with $Y(a^jb^s) = (-1)^s$.

For $r = 1, \cdots, \frac{n-1}{2}$, let $Z^r = \mathbb{C}^2$, $Z^r(a) = \begin{pmatrix} \zeta^r & 0 \\ 0 & \zeta^{-r} \end{pmatrix}$ with $\zeta = e^{2\pi\sqrt{-1}/n}$, and $Z^r(b) = \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}$.

For $j = 1, \cdots, \frac{n-1}{2}$ and $t = 1, \cdots, n$, let $V^{j,t} = \mathbb{C}a^j \oplus \mathbb{C}a^{-j}$, and $V^{j,t}(a) = \begin{pmatrix} \zeta^t & 0 \\ 0 & \zeta^{-t} \end{pmatrix}$, $V^{j,t}(b) = \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}$.

Let $W^{\pm} = \bigoplus_{k=0}^{n-1} \mathbb{C} a^k b$, let $W^{\pm}(a)(a^k b) = a^{k+2} b$, and $W^{\pm}(b)(a^k b) = \pm a^{-k} b$. There is an isomorphism

$$(W^{\pm})^{\star} \cong W^{\pm}, \qquad \sum_{k=0}^{n-1} u_k \cdot a^k b \mapsto \sum_{k=0}^{n-1} \overline{u_k} \cdot a^k b. \tag{99}$$

Then any simple object of $\mathcal{E}$ is isomorphic to one of X, Y, Z^r $(1 \leqslant r \leqslant \frac{n-1}{2})$, $V^{j,t}$ $(1 \leqslant j \leqslant \frac{n-1}{2}, 1 \leqslant t \leqslant n)$, W^+, W^-.

Remark 6.1. *Since* $\chi_U(a^k b, g) = 0$ *unless* $U = W^{\pm}$, *to compute* $CN_n(K)$ *it is enough to compute* $F(K, W^{\pm})$.

Define

$$\beta_{W^{\pm}} : W^{\pm} \odot W^{\pm} \cong X \bigoplus_{r=1}^{\frac{n-1}{2}} Z^r \bigoplus_{j=1}^{\frac{n-1}{2}} \bigoplus_{t=1}^{n} V^{j,t},$$

$$\sum_{k,k'=0}^{n-1} w_{k,k'} \cdot a^k b \otimes a^k b$$

$$\mapsto (\sum_{k=0}^{n-1} w_{k,k}) \bigoplus_{r=1}^{\frac{n-1}{2}} \left(\begin{array}{c} \sum\limits_{k=0}^{n-1} \zeta^{\frac{k}{2} r} w_{k,k} \\ \sum\limits_{k=0}^{n-1} \zeta^{\frac{-k}{2} r} w_{k,k} \end{array} \right) \bigoplus_{j=1}^{\frac{n-1}{2}} \bigoplus_{t=1}^{n} ((\sum_{l=0}^{n-1} \zeta^{\frac{lt}{2}} w_{l+j,l}) a^j + (\sum_{l=0}^{n-1} \zeta^{\frac{-lt}{2}} w_{l-j,l}) a^{-j}, \tag{100}$$

with the inverse

$$\beta_{W^{\pm}}^{-1} \left((x) \bigoplus_{r=1}^{\frac{n-1}{2}} \left(\begin{array}{c} y_r \\ z_r \end{array} \right) \bigoplus_{j=1}^{\frac{n-1}{2}} \bigoplus_{t=1}^{n} (v_{j,t}^{+} a^j + v_{j,t}^{-} a^{-j}) \right) = \sum_{k,k'=0}^{n-1} w_{k,k'} \cdot a^k b \otimes a^k b, \tag{101}$$

where

$$w_{k,k} = \frac{1}{n}(x + \sum_{r=1}^{\frac{n-1}{2}} \zeta^{-\frac{kr}{2}} y_r + \zeta^{\frac{kr}{2}} z_r), \qquad 1 \leqslant k \leqslant n-1, \tag{102}$$

$$w_{l\pm j,l} = \frac{1}{n}(\sum_{t=1}^{n} \zeta^{\mp \frac{lt}{2}} v_{j,t}^{\pm}), \qquad 0 \leqslant l \leqslant n-1, 1 \leqslant j \leqslant \frac{n-1}{2}. \tag{103}$$

6.2 Computing Φ and ROT

Suppose $F : W^{\pm} \odot W^{\pm} \to W^{\pm} \odot W^{\pm}$ is a morphism such that

$$F(a^k b \otimes a^{k'} b) = \sum_{s,s'=0}^{n-1} \mathcal{T}(F)_{k,k'}^{s,s'} \cdot a^s b \otimes a^{s'} b. \tag{104}$$

From $F \circ a = a \circ F$ and $F \circ b = b \circ F$ it follows that

$$T^{s+2,s'+2}_{k+2,k'+2} = T^{s,s'}_{k,k'} = T^{-s,-s'}_{-k,-k'}, \qquad T^{s,s'}_{k,k'} \neq 0 \Rightarrow s - s' = k - k'. \tag{105}$$

Denote $\Phi = \Phi^{W^\pm,W^\pm}_{W^\pm,W^\pm}$. Suppose

$$\Phi(F) = (\lambda)_X \bigoplus_{r=1}^{\frac{n-1}{2}} (\mu_r)_{Z^r} \bigoplus_{j=1}^{\frac{n-1}{2}} \bigoplus_{t=1}^{n} (\nu_{j,t})_{V^{j,t}}, \tag{106}$$

where we use $(\kappa)_V$ to denote $\kappa \cdot \mathrm{id}_V$, thanks to $N^V_{W^\pm,W^\pm} \in \{0,1\}$ for all $V \in \Lambda$.

By direct computation, we get

$$\lambda = \sum_{d=0}^{n-1} \mathcal{T}(F)^{0,0}_{d,d}, \tag{107}$$

$$\mu_r = \sum_{d=0}^{n-1} (\zeta^{\frac{dr}{2}} + \zeta^{-\frac{dr}{2}}) \mathcal{T}(F)^{0,0}_{d,d}, \tag{108}$$

$$\nu_{j,t} = \sum_{d=0}^{n-1} \zeta^{-\frac{dt}{2}} \mathcal{T}(F)^{j,0}_{d+j,d}; \tag{109}$$

the inverse Φ^{-1} appears as

$$\mathcal{T}(F)^{0,0}_{d,d} = \frac{1}{n}(\lambda + \sum_{r=1}^{\frac{n-1}{2}} (\zeta^{\frac{dr}{2}} + \zeta^{-\frac{dr}{2}}) \mu_r), \tag{110}$$

$$\mathcal{T}(F)^{\pm j,0}_{d\pm j,d} = \frac{1}{n} \sum_{t=1}^{n} \zeta^{\frac{\pm dt}{2}} \nu_{j,t}. \tag{111}$$

Since $R_{W^\pm} := R_{W^\pm,W^\pm} : W^\pm \odot W^\pm \to W^\pm \odot W^\pm$ is, by (16), given by

$$\sum_{k,k'=0}^{n-1} w_{k,k'} \cdot (a^k b \otimes a^{k'} b) \mapsto \pm \sum_{k,k'=0}^{n-1} w_{k,k'} \cdot (a^{2k-k'} b \otimes a^k b). \tag{112}$$

By (107)-(109) we conclude that

$$\Phi(R_{W^\pm}) = (\pm 1)_X \bigoplus_{r=1}^{\frac{n-1}{2}} (\pm 1)_{Z^r} \bigoplus_{j=1}^{\frac{n-1}{2}} \bigoplus_{t=1}^{n} (\pm \zeta^{\frac{jt}{2}})_{V^{j,t}}. \tag{113}$$

By (93), (95), and (107)-(111), we have

$$\mathrm{ROT}^{W^\pm} := \mathrm{ROT}^{W^\pm,W^\pm}_{W^\pm,W^\pm} = \begin{pmatrix} \frac{1}{n} & \frac{2}{n} & \frac{2}{n} \\ \frac{1}{n} & \frac{2}{n} & \frac{1}{n}(\zeta^{\frac{jr}{2}} + \zeta^{-\frac{jr}{2}}) \\ \frac{1}{n} & \frac{1}{n}(\zeta^{\frac{jr'}{2}} + \zeta^{-\frac{jr'}{2}}) & \frac{1}{n}(\zeta^{\frac{jt'+j't}{2}} + \zeta^{-\frac{jt'+j't}{2}}) \end{pmatrix} \tag{114}$$

at $\begin{pmatrix} X \\ Z^r \\ V^{j,t} \end{pmatrix} \times (X, Z^{r'}, V^{j',t'})$; by this we mean that

$$\begin{aligned}
(\mathrm{ROT}^{W^{\pm}})_{X,X} = (\mathrm{ROT}^{W^{\pm}})_{Z^r,X} = (\mathrm{ROT}^{W^{\pm}})_{V^{j,t},X} &= \frac{1}{n}, \\
(\mathrm{ROT}^{W^{\pm}})_{X,Z^{r'}} = (\mathrm{ROT}^{W^{\pm}})_{X,V^{j',t'}} = (\mathrm{ROT}^{W^{\pm}})_{Z^r,Z^{r'}} &= \frac{2}{n}, \\
(\mathrm{ROT}^{W^{\pm}})_{Z^r,V^{j,t}} = (\mathrm{ROT}^{W^{\pm}})_{V^{j,t},Z^r} &= \frac{1}{n}(\zeta^{\frac{jr}{2}} + \zeta^{-\frac{jr}{2}}), \\
(\mathrm{ROT}^{W^{\pm}})_{V^{j,t},V^{j',t'}} &= \frac{1}{n}(\zeta^{\frac{jt'+j't}{2}} + \zeta^{-\frac{jt'+j't}{2}}),
\end{aligned}$$

and all the other entries vanish.

6.3 The result

By direct computation, one can obtain

$$\mathrm{ROT}^{W^{\pm}}\Phi(R^c_{W^{\pm}}) = (\pm 1)^c (n,c) \left((1)_X \bigoplus_{r:\ (n,c)|r} (1)_{Z^r} \bigoplus_{\substack{j:\ (n,c)|j \\ 1 \leqslant j \leqslant \frac{n-1}{2}}} \bigoplus_{\substack{t:\ (n,c)|t \\ 1 \leqslant t \leqslant n}} (\zeta^{-\frac{jt}{2c}})_{V^{j,t}} \right).$$

For a rational number $\frac{p}{q} = [[c_1, \cdots, c_k]]$, let

$$\mu(\frac{p}{q}) = \sum_{s=1}^{k} c_s. \tag{115}$$

It can be proved by induction on k that

$$\begin{aligned}
&\check{F}(T(\frac{p}{q}), W^{\pm}) \\
=&(\pm 1)^{\mu(\frac{p}{q})} (n,p) \cdot \left((1)_X \bigoplus_{r:\ (n,p)|r} (1)_{Z^r} \bigoplus_{\substack{j:\ (n,p)|j \\ 1 \leqslant j \leqslant \frac{n-1}{2}}} \bigoplus_{\substack{t:\ (n,p)|t \\ 1 \leqslant t \leqslant n}} (\zeta^{-\frac{q}{2p}jt})_{V^{j,t}} \right).
\end{aligned} \tag{116}$$

Using the following two number-theoretical facts

- $[(n,p_1), \cdots, (n,p_m)] = (n, [p_1, \cdots, p_m])$,
- $\sum\limits_{\substack{j:\ (n,A)|j \\ 1 \leqslant j \leqslant \frac{n-1}{2}}} \bigoplus\limits_{\substack{t:\ (n,A)|t \\ 1 \leqslant t \leqslant n}} \zeta^{-\frac{B}{2A}jt} = \frac{n}{2(n,A)} \cdot ((\frac{n}{(n,A)}, B) - 1)$ for any $A, B \in \mathbb{Z} - \{0\}$,

one can further deduce that

$$\begin{aligned}
&|\overrightarrow{Tr}(\check{F}(T(\frac{p_m}{q_m}), W^{\pm}) \circ \cdots \circ \check{F}(T(\frac{p_1}{q_1}), W^{\pm}))| \\
=&(\pm 1)^{\sum_{i=1}^{m} \mu(\frac{p_i}{q_i})} n \prod_{i=1}^{m} (n,p_i) \cdot \frac{(n/(n,[p_1,\cdots,p_m]), N(\frac{p_1}{q_1}, \cdots, \frac{p_m}{q_m}))}{(n,[p_1,\cdots,p_m])},
\end{aligned} \tag{117}$$

with

$$N(\frac{p_1}{q_1},\cdots,\frac{p_m}{q_m})=[p_1,\cdots,p_m]\cdot\sum_{i=1}^{m}\frac{q_i}{p_i}. \tag{118}$$

Finally, combining Theorem 5.4, Remark 6.1, (7), (96), (98), (117), and noting that $\theta_{W^\pm}=\pm1$ and $\sum\limits_{i=1}^{m}\mu(\frac{p_i}{q_i})\equiv\mathrm{wr}(M(\frac{p_1}{q_1},\cdots,\frac{p_m}{q_m}))\pmod 2$, we obtain

Theorem 6.2. *The number of n-colorings of the Montesinos knot $M(\frac{p_1}{q_1},\cdots,\frac{p_m}{q_m})$ is*

$$n\prod_{i=1}^{m}(n,p_i)\cdot\frac{(n/(n,[p_1,\cdots,p_m]),N(\frac{p_1}{q_1},\cdots,\frac{p_m}{q_m}))}{(n,[p_1,\cdots,p_m])}.$$

In particular, the number of n-colorings of the rational knot $M(\frac{p}{q})$ is $n(n,q)$, and that of the pretzel knot $M(p_1,\cdots,p_m)$ is

$$n\prod_{i=1}^{m}(n,p_i)\cdot\frac{(n/(n,[p_1,\cdots,p_m]),N(p_1,\cdots,p_m))}{(n,[p_1,\cdots,p_m])}.$$

7 Appendix

7.1 Proof of Theorem 3.5

Proof. (a) The inverse f^{-1} induces an isomorphism of fundamental groupoids $(f^{-1})_*:\Pi_1(B')\to\Pi_1(B)$; it further induces an isomorphism of categories $\mathfrak{B}^\Gamma(B)\to\mathfrak{B}^\Gamma(B')$, and then the desired isomorphism $f_*:Z(B)\to Z(B')$.

(b) Obviously.

(c) Suppose $\#\pi_0(\partial B)=l$ and $\#\pi_0(A)=k$. By definition, for $\mathbf{y}\in\Gamma^l$,

$$(\mathrm{gl}_A(Z(B^{\mathrm{cut}})))_{\mathbf{y}}=\frac{1}{(\#\Gamma)^k}\cdot\left(\bigoplus_{\mathbf{x}\in\Gamma^k}(\bigoplus_{[\rho]\in\pi_0(\mathfrak{B}^\Gamma_{\mathbf{y},\mathbf{x},\mathbf{x^{-1}}})}\mathbb{C}[\rho])\right)^{\Gamma^k}.$$

Define

$$\Theta_{\mathbf{y}}:(\mathrm{gl}_A(Z(B^{\mathrm{cut}})))_{\mathbf{y}}\to Z(B)_{\mathbf{y}}=\bigoplus_{[\tau]\in\mathfrak{B}^\Gamma_{\mathbf{y}}(B)}\mathbb{C}[\tau]$$

by sending each Γ^k-invariant element $\sum\limits_{\mathbf{x}\in\Gamma^k}\sum\limits_{[\rho]\in\pi_0(\mathfrak{B}^\Gamma_{\mathbf{y},\mathbf{x},\mathbf{x^{-1}}})}\lambda_{\mathbf{x},[\rho]}\cdot[\rho]$ to $\frac{1}{(\#\Gamma)^k}\cdot\sum\limits_{\mathbf{x}\in\Gamma^k}\sum\limits_{[\rho]\in\pi_0(\mathfrak{B}^\Gamma_{\mathbf{y},\mathbf{x},\mathbf{x^{-1}}})}\lambda_{\mathbf{x},[\rho]}\cdot[\mathrm{gl}(\rho)]$, where $\mathrm{gl}(\rho)\in\mathfrak{B}^\Gamma(B)$ is obtained by "gluing ρ along A. It is easy to see that $\Theta_{\mathbf{y}}$ is well-defined, and these $\Theta_{\mathbf{y}}$ together define a morphism $\Theta:\mathrm{gl}_A(Z(B^{\mathrm{cut}}))\to Z(B)$.

Define

$$\Psi_{\mathbf{y}} : Z(B)_{\mathbf{y}} \to (\mathrm{gl}_A(Z(B^{\mathrm{cut}})))_{\mathbf{y}}$$

by sending $[\tau]$ to $\sum_{\mathbf{g}\in\Gamma^k} \eta_{\mathbf{g},\mathbf{g}} \cdot [\tau]$. These $\Psi_{\mathbf{y}}$ together define a morphism $\Psi : Z(B) \to \mathrm{gl}_A(Z(B^{\mathrm{cut}}))$, and it can be checked that $\Psi = \Theta^{-1}$. ☐

7.2 Proof of Theorem 3.13

Proof. (a) and (b) are obvious.

(c)-(i) $\hat{Z}(B'_{v,2})(Z(C))_U$ is induced by

$$\begin{aligned}
&\sum_{\mathbf{h},\mathbf{g}} \delta(\eta_{\mathbf{e},\mathbf{h}} \cdot [\psi'_2]) \otimes \delta(\eta_{\mathbf{h},\mathbf{g}} \cdot [\psi_2]) \otimes \delta(\eta_{\mathbf{g},\mathbf{e}} \cdot [\phi]) \\
&\mapsto \sum_{\substack{\mathbf{h},\mathbf{g}',\mathbf{y}' \\ [\phi'],[\psi_1]}} Y(C)^{[\phi'],[\psi_1]}_{[\psi_2],[\phi]} \cdot \delta(\eta_{\mathbf{e},\mathbf{h}} \cdot [\psi'_2]) \otimes \delta(\eta_{\mathbf{h},\mathbf{g}'} \cdot [\phi']) \otimes \delta(\eta_{\mathbf{g}',\mathbf{e}} \cdot [\psi_1]),
\end{aligned}$$

and $Z(C')_{\hat{Z}(B_{v,1})(U)}$ is induced by

$$\begin{aligned}
&\sum_{\mathbf{h},\mathbf{g}'} \delta(\eta_{\mathbf{e},\mathbf{h}} \cdot [\psi'_2]) \otimes \delta(\eta_{\mathbf{h},\mathbf{g}'} \cdot [\phi']) \otimes \delta(\eta_{\mathbf{g}',\mathbf{e}} \cdot [\psi_1]) \\
&\mapsto \sum_{\substack{\mathbf{h}',\mathbf{g}',\mathbf{z}' \\ [\phi''],[\psi'_1]}} Y(C')^{[\phi''],[\psi'_1]}_{[\phi'],[\psi_1]} \cdot \delta(\eta_{\mathbf{e},\mathbf{h}'} \cdot [\phi'']) \otimes \delta(\eta_{\mathbf{h}',\mathbf{g}'} \cdot [\psi'_1]) \otimes \delta(\eta_{\mathbf{g}',\mathbf{e}} \cdot [\psi_1]).
\end{aligned}$$

Hence $Z(C')_{\hat{Z}(B_{v,1})(U)} \circ \hat{Z}(B'_{v,2})(Z(C))_U$ is induced by

$$\begin{aligned}
&\sum_{\mathbf{h},\mathbf{g}} \delta(\eta_{\mathbf{e},\mathbf{h}} \cdot [\psi'_2]) \otimes \delta(\eta_{\mathbf{h},\mathbf{g}} \cdot [\psi_2]) \otimes \delta(\eta_{\mathbf{g},\mathbf{e}} \cdot [\phi]) \\
\mapsto &\sum_{\substack{\mathbf{h}',\mathbf{g}',\mathbf{y}',\mathbf{z}' \\ [\phi''],[\psi'_1],[\psi_1]}} \left(\sum_{[\phi']} Y(C)^{[\phi'],[\psi_1]}_{[\psi_2],[\phi]} \cdot Y(C')^{[\phi''],[\psi'_1]}_{[\phi'],[\psi_1]} \right) \\
&\cdot \delta(\eta_{\mathbf{e},\mathbf{h}'} \cdot [\phi'']) \otimes \delta(\eta_{\mathbf{h}',\mathbf{g}'} \cdot [\psi'_1]) \otimes \delta(\eta_{\mathbf{g}',\mathbf{e}} \cdot [\psi_1]) \\
= &\sum_{\substack{\mathbf{h}',\mathbf{g}',\mathbf{y}',\mathbf{z}' \\ [\phi''],[\psi'_1],[\psi_1]}} Y(C_v)^{[\phi''],[\psi'_1 * \psi_1]}_{[\psi'_2 * \psi_2],[\phi]} \cdot \delta(\eta_{\mathbf{h}',\mathbf{e}} \cdot [\phi'']) \otimes \delta(\eta_{\mathbf{h}',\mathbf{g}'} \cdot [\psi'_1]) \otimes \delta(\eta_{\mathbf{g}',\mathbf{e}} \cdot [\psi_1]),
\end{aligned}$$

which is just the map inducing $Z(C_v)_U$. In the last line $\psi'_1 * \psi_1$ means the morphism obtained by "gluing" ψ'_1 and ψ_1, and we have used the key fact that

$$\sum_{[\phi']} Y(C)^{[\phi'],[\psi_1]}_{[\psi_2],[\phi]} \cdot Y(C')^{[\phi''],[\psi'_1]}_{[\phi'],[\psi_1]} = Y(C_v)^{[\phi''],[\psi'_1 * \psi_1]}_{[\psi'_2 * \psi_2],[\phi]}.$$

(c)-(ii) The proof is similar to (c)-(i). ☐

7.3 Proof of Proposition 3.14

Proof. (a) $Z(C')_{\alpha'} \circ ((F|_{B_{h,1}})_*)_U$

$$
\begin{aligned}
&=\hat{Z}(B'_{h,2})(\alpha \circ ((F^{-1}|_{B'_{v,1}})_*)_U) \circ Z(C')_U \circ ((F|_{B_{h,1}})_*)_U \\
&=\hat{Z}(B'_{h,2})(\alpha \circ ((F^{-1}|_{B'_{v,1}})_*)_U) \circ ((\partial F)_*)_U \circ Z(C)_U \qquad (119)\\
&=\hat{Z}(B'_{h,2})(\alpha) \circ \hat{Z}(B'_{h,2})(((F^{-1}|_{B'_{v,1}})_*)_U) \circ \\
&\quad \hat{Z}(B'_{h,2})(((F|_{B_{v,1}})_*)_U) \circ ((F|_{B_{h,2}})_*)_{\hat{Z}(B_{v,1})(U)} \circ Z(C)_U \\
&=\hat{Z}(B'_{h,2})(\alpha) \circ ((F|_{B_{h,2}})_*)_{\hat{Z}(B_{v,1})(U)} \circ Z(C)_U \\
&=((F|_{B_{h,2}})_*)_{U'} \circ \hat{Z}(B_{h,2})(\alpha) \circ Z(C)_U \qquad (120)\\
&=((F|_{B_{h,2}})_*)_{U'} \circ Z(C)_\alpha,
\end{aligned}
$$

where in (119) $(\partial F)_*$ means the natural transformation $\hat{Z}(B_{h,2}) \circ \hat{Z}(B_{v,1}) \to \hat{Z}(B'_{h,2}) \circ \hat{Z}(B'_{v,1})$ and Theorem 3.13 (a) is applied, and in (120) we have applied the naturality of $(F|_{B_{h,2}})_*$ to α.

(b) $Z(C_v)_{\alpha' \circ \hat{Z}(B'_{v,1})(\alpha)}$

$$
\begin{aligned}
&=\hat{Z}(B'_{h,2})(\alpha' \circ \hat{Z}(B'_{v,1})(\alpha)) \circ Z(C_v)_U \\
&=\hat{Z}(B'_{h,2})(\alpha') \circ \hat{Z}(B'_{h,2})\hat{Z}(B'_{v,1})(\alpha) \circ Z(C')_{\hat{Z}(B_{v,1})(U)} \circ Z(C)_U \\
&=\hat{Z}(B'_{h,2})(\alpha') \circ Z(C')_{U'} \circ \hat{Z}(B_{h,2})(\alpha) \circ Z(C)_U \qquad (121)\\
&=Z(C')_{\alpha'} \circ Z(C)_\alpha,
\end{aligned}
$$

where in (121) we have applied the naturality of $Z(C')$ to α.

(c) $Z(C^1 \odot C^2)_{\alpha^1 \boxtimes \alpha^2}$

$$
\begin{aligned}
&=\hat{Z}(B_{h,2}(\mathbf{0}))(\hat{Z}(B^1_{h,2})(\alpha^1) \boxtimes \hat{Z}(B^2_{h,2})(\alpha^2)) \circ Z(C^1 \odot C^2)_{U^1 \boxtimes U^2} \\
&=(\hat{Z}(B^1_{h,2})(\alpha^1) \odot \hat{Z}(B^2_{h,2})(\alpha^2)) \circ (Z(C^1)_{U^1} \odot Z(C^2)_{U^2}) \qquad (122)\\
&=(\hat{Z}(B^1_{h,2})(\alpha^1) \circ Z(C^1)_{U^1}) \odot (\hat{Z}(B^2_{h,2})(\alpha^2) \circ Z(C^2)_{U^2}) \\
&=Z(C^1)_{\alpha^1} \odot Z(C^2)_{\alpha^2},
\end{aligned}
$$

where in (122) we apply Theorem 3.5 (c)-(ii) to the case $C = C^1 \sqcup C^2, C' = C(\mathbf{0})$, using $\hat{Z}(B_{h,2}(\mathbf{0})) \cong \odot$ and $Z(C(\mathbf{0}))_{\hat{Z}(B^1_{h,1})(U^1) \boxtimes \hat{Z}(B^2_{h,1})(U^2)} = \mathrm{id}$. □

References

[1] D.Altschuler, A.Coste, *Quasi-quantum groups, knots, three-manifolds, and topological field theory*. Communications in Mathematical Physics, 150-1 (1992) 83-107.

[2] B.Bakalov, *Lectures on tensor categories and modular functors*. In: University Lecture Series, vol.21, American Mathematical Society, Providence, Rhode Island, USA, 2001.

[3] H.M.Chen, *Applying TQFT to count regular coverings of Seifert 3-manifolds*. Journal of Geometry and Physics 62-6 (2012) 1347-1357.

[4] R.H.Crowell, R.H.Fox, *Introduction to knot theory*. Graduate Texts in Mathematics, vol.57, Springer-Verlag, New York Heidelberg Berlin, 1963.

[5] R.Dijkgraaf, E.Witten, *Topological gauge theories and group cohomology*. Communications in Mathematical Physics 129-2 (1990) 393-429.

[6] M.Eisermann, *Knot colouring polynomials*. Pacific Journal of Mathematic 231-2 (2007) 305-336.

[7] K.Ferguson, *Link invariants associated to TQFT's with finite gauge groups*. Journal of Knot Theory and Its Ramification 2-1 (1993) 11-36.

[8] D.S.Freed, Quinn, *Chern-Simons theory with finite gauge group*. Communications in Mathematical Physics 156-3 (1993) 435-472.

[9] D.S.Freed, *Higher algebraic structures and quantization*. Communications in Mathematical Physics 159-2 (1994) 343-398.

[10] P.Hall, *The Eulerian functions of a group*. The Quarterly Journal of Mathematics 7 (1936) 134-151.

[11] G.Jones, *Enumeration of homomorphisms and surface-coverings*. The Quarterly Journal of Mathematics 46-2 (1995) 485-507.

[12] G.Laures, *On cobordism of manifolds with corners*. Transactions of the american mathematical society 352-12 (2000) 5667-5688.

[13] V.Liskovets, A.Mednykh, *Enumeration of subgroups in the fundamental groups of orientable circle bundles over surfaces*. Communications in Algebra 28-4 (2000) 1717-1738.

[14] A.Lubotzky, *Counting finite index subgroups*. In: Groups '93, Galway/St.Andrews, Vol.2, London Mathametical Society, Lecture Note Series, vol.212, Cambridge University Press, Cambridge, 1995, 368-404.

[15] D.Matei, A.I.Suciu, *Hall invariants, homology of subgroups, and characteristic varieties*. International Mathematics Research Notices 9 (2002) 465-503.

[16] J.C.Morton, *Extended TQFT, gauge theory, and 2-linearization*. arXiv:1003.5603v3.

[17] T.Ohtsuki, *Problems on invariants of knots and 3-manifolds*. Geometry & Topology Monographs, Vol.4 (2002)

[18] E.Pervova, *On the existence of branced coverings between surfaces with prescribed branch data I*. Algebraic & Geometric Topology 6 (2006) 1957-1985.

[19] J.H.Przytycki, *3-coloring and other elementary invariants of knots*. In: “Knot theory”, Banach Center Publ.42, Polish Acad. Sci. Warsaw (1998) 275-295.

[20] N.Reshetikhin, V.G.Turaev, *Invariants of 3-manifolds via link polynomials and quantum groups*. Inventiones Mathematicae 103 (1991) 547-597.

[21] D.S.Silver, S.G.Williams, *Generalized n-colorings of links*. In: “Knot theory”, Banach Center Publ.42, Polish Acad. Sci. Warsaw (1998) 381-394.

[22] A.I.Suciu, *Fundamental groups of line arrangements: Enumerative aspects*. In: Advances in algebraic geometry motivated by physics (Lowell, MA, 2000), 43C79, Contemp. Math., 276, Amer. Math. Soc., Providence, RI, 2001.

[23] V.G.Turaev, *On certain enumeration problems in two-dimensional topology*. Mathametical Research Letter 16-3 (2009) 515-529.

[24] M.Wakui, *On Dijkgraaf-Witten invariant for 3-manifolds*. Osaka Journal of Mathematics 29 (1992) 675-696.